

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PegIntron safely and effectively. See full prescribing information for PegIntron.

PegIntron (peginterferon alfa-2b) Injection, Powder for Solution for Subcutaneous Use

Initial U.S. Approval: 2001

WARNING: RISK OF SERIOUS DISORDERS AND RIBAVIRIN-ASSOCIATED EFFECTS

See full prescribing information for complete boxed warning.

- May cause or aggravate fatal or life-threatening neuropsychiatric, autoimmune, ischemic, and infectious disorders. Monitor closely and withdraw therapy with persistently severe or worsening signs or symptoms of the above disorders (5)

Use with Ribavirin

- Ribavirin may cause birth defects and fetal death; avoid pregnancy in female patients and female partners of male patients (5.1)
- Ribavirin is a potential carcinogen (5.1, 13.1)

RECENT MAJOR CHANGES

- Indications and Usage, Chronic Hepatitis C (1.1) [12/2008]
- Dosage and Administration, Combination Therapy (2.1) [12/2008]
- Dosage and Administration, Dose Reduction (2.3) [12/2008]
- Dosage and Administration, Discontinuation of Dosing (2.4) [12/2008]
- Contraindications (4) [12/2008]
- Warnings and Precautions, Laboratory Tests (5.15) [12/2008]
- Warnings and Precautions, Impact on Growth-Pediatric Use (5.18) [12/2008]

INDICATIONS AND USAGE

PegIntron is an antiviral indicated for

- **Combination therapy** with REBETOL (ribavirin): Chronic Hepatitis C (CHC) in patients ≥3 years with compensated liver disease previously untreated with interferon alpha (1.1)
- **Monotherapy**: CHC in patients (≥18 years) with compensated liver disease previously untreated with interferon alpha (1.1)

DOSAGE AND ADMINISTRATION

- PegIntron is administered by subcutaneous injection

	PegIntron Dose (Adults) ^a	PegIntron Dose (Pediatric Patients)	REBETOL Dose ^a (Adults)	REBETOL Dose (Pediatric Patients)
PegIntron/REBETOL Combination Therapy (2.1)	1.5 mcg/kg/week	60 mcg/m ² /week	800-1400 mg orally daily with food	15 mg/kg/day orally with food in two divided doses

a-Refer to Tables 1 and 3 of the full Prescribing Information.

- Dose reduction is recommended in patients experiencing certain adverse reactions or renal dysfunction (2.3, 2.5)

DOSAGE FORMS AND STRENGTHS

Single-use vial (with 1.25 mL diluent) and REDIPEN (3):

- 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL, 150 mcg per 0.5 mL

CONTRAINDICATIONS

- Known hypersensitivity reactions, such as urticaria, angioedema, bronchoconstriction, anaphylaxis, Stevens Johnson syndrome, and toxic epidermal necrolysis to interferon alpha or any other product component (4)
 - Autoimmune hepatitis (4)
 - Hepatic decompensation (Child-Pugh score >6 [class B and C]) in cirrhotic CHC patients before or during treatment (4)
- Additional contraindications for combination therapy with ribavirin:

- Pregnant women and men whose female partners are pregnant (4, 8.1)
- Hemoglobinopathies (e.g., thalassemia major, sickle-cell anemia) (4)
- Creatinine clearance <50 mL/min (4)

WARNINGS AND PRECAUTIONS

- Birth defects and fetal death with ribavirin: Patients must have a negative pregnancy test prior to therapy; use at least 2 forms of contraception and undergo monthly pregnancy tests (5.1)

Patients exhibiting the following conditions should be closely monitored and may require dose reduction or discontinuation of therapy:

- Hemolytic anemia with ribavirin (5.1)
- Neuropsychiatric events (5.2)
- History of significant or unstable cardiac disease (5.3)
- Hypothyroidism, hyperthyroidism, hyperglycemia, diabetes mellitus that cannot be effectively treated by medication (5.4)
- New or worsening ophthalmologic disorders (5.5)
- Ischemic and hemorrhagic cerebrovascular events (5.6)
- Severe decreases in neutrophil or platelet counts (5.7)
- History of autoimmune disorders (5.8)
- Pancreatitis and ulcerative or hemorrhagic/ischemic colitis and pancreatitis (5.9, 5.10)
- Pulmonary infiltrates or pulmonary function impairment (5.11)
- Child-Pugh score >6 (class B and C) (4, 5.12)
- Increased creatinine levels in patients with renal insufficiency (5.13)
- Serious, acute hypersensitivity reactions and cutaneous eruptions (5.14)
- Dental/periodontal disorders reported with combination therapy (5.16)
- Hypertriglyceridemia may result in pancreatitis (e.g., triglycerides >1000 mg/dL) (5.17)
- Weight loss and growth inhibition reported with combination therapy in pediatric patients (5.18)

ADVERSE REACTIONS

Most common adverse reactions (>40%) in adult patients receiving either PegIntron or PegIntron/REBETOL are injection site inflammation/reaction, fatigue/asthenia, headache, rigors, fevers, nausea, myalgia and anxiety/emotional lability/irritability (6.1). Most common adverse reactions (>25%) in pediatric patients receiving PegIntron/REBETOL are pyrexia, headache, neutropenia, fatigue, anorexia, injection site erythema, vomiting (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Schering Corporation at 1-800-526-4099 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Drug Metabolized by CYP450: Caution with drugs metabolized by CYP2C8/9 (e.g., warfarin, phenytoin) or CYP2D6 (e.g., flecainide) (7.1)
- Methadone: Monitor for increased narcotic effect (7.2)
- Nucleoside Analogues: Closely monitor for toxicities. Discontinue Nucleoside Reverse Transcriptase Inhibitors or reduce dose or discontinue interferon, ribavirin, or both with worsening toxicities (7.3)
- Didanosine: Concurrent use with REBETOL is not recommended (7.3)

USE IN SPECIFIC POPULATIONS

- Ribavirin Pregnancy Registry: 1-800-593-2214 (8.1)
- Pediatrics: safety and efficacy in pediatrics <3 years old have not been established (8.4)
- Geriatrics: neuropsychiatric, cardiac, pulmonary, GI, and systemic (flu-like) adverse reactions may be more severe (8.5)
- Organ transplant: safety and efficacy have not been studied (8.6)
- HIV or HBV co-infection: safety and efficacy have not been established (8.7)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: [12/2008]

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING – RISK OF SERIOUS DISORDERS AND RIBAVIRIN-ASSOCIATED EFFECTS

1 INDICATIONS AND USAGE

1.1 Chronic Hepatitis C

2 DOSAGE AND ADMINISTRATION

- 2.1 PegIntron/REBETOL Combination Therapy
- 2.2 PegIntron Monotherapy
- 2.3 Dose Reduction
- 2.4 Discontinuation of Dosing
- 2.5 Renal Function
- 2.6 Preparation and Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Use with Ribavirin
- 5.2 Neuropsychiatric Events
- 5.3 Cardiovascular Events
- 5.4 Endocrine Disorders
- 5.5 Ophthalmologic Disorders
- 5.6 Cerebrovascular Disorders
- 5.7 Bone Marrow Toxicity
- 5.8 Autoimmune Disorders
- 5.9 Pancreatitis
- 5.10 Colitis
- 5.11 Pulmonary Disorders
- 5.12 Hepatic Failure
- 5.13 Patients with Renal Insufficiency
- 5.14 Hypersensitivity
- 5.15 Laboratory Tests
- 5.16 Dental and Periodontal Disorders
- 5.17 Triglycerides
- 5.18 Impact on Growth- Pediatric Use

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Immunogenicity
- 6.3 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drugs Metabolized by Cytochrome P-450
- 7.2 Methadone
- 7.3 Use with Ribavirin (Nucleoside Analogues)

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Organ Transplant Recipients
- 8.7 HIV or HBV Coinfection

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Chronic Hepatitis C in Adults
- 14.2 Chronic Hepatitis C in Pediatrics

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Medication Guide
- 17.2 Pregnancy
- 17.3 HCV Transmission
- 17.4 Laboratory Evaluations, Hydration, “Flu-like” Symptoms

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: RISK OF SERIOUS DISORDERS AND RIBAVIRIN-ASSOCIATED EFFECTS

Alpha interferons, including PegIntron, may cause or aggravate fatal or life-threatening neuropsychiatric, autoimmune, ischemic, and infectious disorders. Patients should be monitored closely with periodic clinical and laboratory evaluations. Patients with persistently severe or worsening signs or symptoms of these conditions should be withdrawn from therapy. In many, but not all cases, these disorders resolve after stopping PegIntron therapy. [See Warnings and Precautions (5) and Adverse Reactions (6.1)]

Use with Ribavirin

Ribavirin may cause birth defects and death of the unborn child. Extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients. Ribavirin causes hemolytic anemia. The anemia associated with REBETOL therapy may result in a worsening of cardiac disease. Ribavirin is genotoxic and mutagenic and should be considered a potential carcinogen. [See REBETOL package insert]

1 INDICATIONS AND USAGE

1.1 Chronic Hepatitis C

Combination therapy:

PegIntron™ in combination with REBETOL® (ribavirin) is indicated for the treatment of chronic hepatitis C in patients 3 years of age and older with compensated liver disease previously untreated with interferon alpha.

Monotherapy (for patients who are intolerant to ribavirin):

PegIntron (peginterferon alfa-2b) is indicated for use alone for the treatment of chronic hepatitis C in patients with compensated liver disease previously untreated with interferon alpha and who are at least 18 years of age.

The following points should be considered when initiating therapy with PegIntron:

- These indications are based on achieving undetectable HCV RNA after treatment for 24 or 48 weeks and maintaining a Sustained Virologic Response (SVR) 24 weeks after the last dose.
- For treatment of CHC, combination therapy with REBETOL is preferred over PegIntron monotherapy unless there are contraindications to or significant intolerance of REBETOL. Combination therapy provides substantially better response rates than monotherapy. [see Clinical Studies (14)]
- No safety and efficacy data are available for treatment of longer than one year.
- The safety and efficacy of PegIntron, alone or in combination with REBETOL, have not been established in patients who have failed other alpha interferon treatments.

2 DOSAGE AND ADMINISTRATION

2.1 PegIntron/REBETOL Combination Therapy

The treatment duration for patients with genotype 1 is 48 weeks. Patients with genotype 2 and 3 should be treated for 24 weeks. REBETOL should be taken with food. REBETOL should not be used in patients with creatinine clearance <50 mL/min.

Adults

The recommended dose of PegIntron is 1.5 mcg/kg/week subcutaneously in combination with 800-1400 mg of REBETOL orally based on patient body weight. The volume of PegIntron to be injected depends on the strength of PegIntron and patient's body weight [See Table 1].

**Table 1
Recommended PegIntron Combination Therapy Dosing (Adults)**

Body weight kg (lbs)	PegIntron REDIPEN® or Vial Strength to Use	Amount of PegIntron (mcg) to Administer	Volume (mL)* of PegIntron to Administer	REBETOL Daily Dose	REBETOL Number of Capsules
<40 (<87)	50 mcg per 0.5 mL	50	0.5	800 mg/day	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
40 – 50 (87 – 111)	80 mcg per 0.5 mL	64	0.4	800 mg/day	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
51 – 60 (112 – 133)		80	0.5	800 mg/day	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
61 – 65 (134 – 144)	120 mcg per 0.5 mL	96	0.4	800 mg/day	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
66 – 75 (145 – 166)		96	0.4	1000 mg/day	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
76 – 85 (167 – 188)		120	0.5	1000 mg/day	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
86 – 105 (189 – 231)	150 mcg per 0.5 mL	150	0.5	1200 mg/day	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
>105 (>231)				1400 mg/day	3 x 200 mg capsules A.M. 4 x 200 mg capsules P.M.

* When reconstituted as directed.

Pediatric Patients

Dosing for pediatric patients is determined by body surface area for PegIntron and by body weight for REBETOL. The recommended dose of PegIntron is 60mcg/m²/week subcutaneously in combination with 15 mg/kg/day of REBETOL orally in two divided doses [see Table 2] for pediatric patients ages 3-17 years. Patients who reach their 18th birthday while receiving PegIntron/REBETOL, should remain on the pediatric dosing regimen.

Table 2
Recommended REBETOL® Dosing in Combination Therapy (Pediatrics)

Body weight kg (lbs)	REBETOL Daily Dose	REBETOL Number of Capsules
<47 (<103)	15 mg/kg/day	Use REBETOL Oral Solution **
47 – 59 (103-131)	800 mg/day	2 x 200 mg capsules A.M. 2 x 200 mg capsules P.M.
60 – 73 (132-162)	1000 mg/day	2 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.
>73 (>162)	1200 mg/day	3 x 200 mg capsules A.M. 3 x 200 mg capsules P.M.

REBETOL to be used in combination with PegIntron 60 mcg/m² weekly.

** REBETOL Oral Solution may be used for any patient regardless of body weight.

2.2 PegIntron Monotherapy

The recommended dose of PegIntron regimen is 1 mcg/kg/week subcutaneously for one year administered on the same day of the week. The volume of PegIntron to be injected depends on patient weight [see Table 3].

Table 3
Recommended PegIntron Monotherapy Dosing

Body weight kg (lbs)	PegIntron REDIPEN or Vial Strength to Use	Amount of PegIntron (mcg) to Administer	Volume (mL)* of PegIntron to Administer
≤45 (≤100)	50 mcg per 0.5 mL	40	0.4
46 – 56 (101 – 124)		50	0.5
57 – 72 (125 – 159)	80 mcg per 0.5 mL	64	0.4
73 – 88 (160 – 195)		80	0.5
89 – 106 (196 – 234)	120 mcg per 0.5 mL	96	0.4
107 – 136 (235 – 300)		120	0.5
137 – 160 (301 – 353)	150 mcg per 0.5 mL	150	0.5

* When reconstituted as directed.

2.3 Dose Reduction

If a serious adverse reaction develops during the course of treatment [see Warning and Precautions (5)] discontinue or modify the dosage of PegIntron and REBETOL until the adverse event abates or decreases in severity. If persistent or recurrent serious adverse events develop despite adequate dosage adjustment, discontinue treatment. For guidelines for dose modifications and discontinuation based on laboratory parameters, see Tables 4 and 5. Dose reduction of PegIntron in adults may be accomplished by utilizing a lower dose strength as shown in Table 6 or 7. For vials, 50% dose reduction may also be accomplished by reducing the volume administered by one-half without changing the dose strength.

In the adult combination therapy trial dose reductions occurred in 42% of subjects receiving PegIntron 1.5 mcg/kg plus REBETOL 800 mg daily including 57% of those subjects weighing 60 kg or less [see Adverse Reactions (6.1)].

Dose reduction in pediatric patients is accomplished by modifying the recommended dose in a two-step process from the original starting dose of 60 mcg/m²/week, to 40 mcg/m²/week, then to 20 mcg/m²/week, if needed [see Tables 4 and 5]. In the pediatric combination therapy trial, dose reductions occurred in 25% of subjects receiving PegIntron 60 mcg/m² weekly plus REBETOL 15 mg/kg daily.

Table 4
Guidelines for Modification or Discontinuation of PegIntron or PegIntron/REBETOL and for Scheduling Visits for Patients with Depression

Depression Severity ¹	Initial Management (4-8 weeks)			Depression	
	Dose modification	Visit schedule	Remains stable	Improves	Worsens
Mild	No change	Evaluate once weekly by visit or phone.	Continue weekly visit schedule.	Resume normal visit schedule.	See moderate or severe depression

Moderate	Adults: Decrease IFN dose 50% Pediatrics: Decrease dose to 40 mcg/m ² /week, then to 20 mcg/m ² /week, if needed	Evaluate once weekly (office visit at least every other week).	Consider psychiatric consultation. Continue reduced dosing.	If symptoms improve and are stable for 4 weeks, may resume normal visit schedule. Continue reduced dosing or return to normal dose.	See severe depression
Severe	Discontinue IFN/R permanently.	Obtain immediate psychiatric consultation.	Psychiatric therapy as necessary		

¹ See DSM-IV for definitions.

Table 5.
Guidelines for Dose Modification and Discontinuation of PegIntron or PegIntron/REBETOL Based on Laboratory Parameters in Adults and Pediatrics

Laboratory Values	PegIntron		REBETOL	
	Adults	Pediatrics	Adults	Pediatrics
Hgb < 10g/dL	For patients with cardiac disease, reduce by 50%*	See footnote*	Decrease by 200mg/day	1 st reduction to 12mg/kg/day 2 nd reduction to 8mg/kg/day
WBC < 1.5 x 10 ⁹ /L Neutrophils < 0.75 x 10 ⁹ /L Platelets < 80 x 10 ⁹ /L (Adults) < 70 x 10 ⁹ /L (Pediatrics)	Reduce by 50%	1 st reduction to 40mcg/m ² /week 2 nd reduction to 20mcg/m ² /week	No Dose Change	No Dose Change
Hgb < 8.5g/dL WBC < 1 x 10 ⁹ /L Neutrophils < 0.5 x 10 ⁹ /L Platelets < 50 x 10 ⁹ /L Creatinine > 2 mg/dL (Pediatrics)	Permanently Discontinue	Permanently Discontinue	Permanently Discontinue	Permanently Discontinue

* For adult patients with a history of stable cardiac disease receiving PegIntron in combination with ribavirin, the PegIntron dose should be reduced by half and the ribavirin dose by 200 mg/day if a >2 g/dL decrease in hemoglobin is observed during any 4-week period. Both PegIntron and ribavirin should be permanently discontinued if patients have hemoglobin levels <12 g/dL after this ribavirin dose reduction. Pediatric patients who have pre-existing cardiac conditions and experience a hemoglobin decrease ≥2 g/dL during any 4-week period during treatment should have weekly evaluations and hematology testing.

Table 6
Reduced PegIntron Dose (0.5 mcg/kg) for (1 mcg/kg) Monotherapy in Adults

Body weight kg(lbs)	PegIntron REDIPEN/Vial Strength to Use	Amount of PegIntron (mcg) to Administer	Volume (mL)** of PegIntron to Administer
≤45 (≤100)	50 mcg per 0.5 mL*	20	0.2
46 – 56 (101 – 124)		25	0.25
57 – 72 (125 – 159)	50 mcg per 0.5 mL	30	0.3
73 – 88 (160 – 195)		40	0.4
89 – 106 (196 – 234)	50 mcg per 0.5 mL	50	0.5
107 – 136 (235 – 300)	80 mcg per 0.5 mL	64	0.4
137 – 160 (301 – 353)		80	0.5

* Must use vial. Minimum delivery for REDIPEN 0.3 mL.

** When reconstituted as directed.

Table 7
Reduced PegIntron Dose (0.75 mcg/kg) for (1.5 mcg/kg) Combination Therapy in Adults

Body weight kg(lbs)	PegIntron REDIPEN/Vial Strength to Use	Amount of PegIntron (mcg) to Administer	Volume (mL)** of PegIntron to Administer
<40 (<87)	50 mcg per 0.5 mL*	25	0.25
40 – 50 (87 – 111)	50 mcg per 0.5 mL	30	0.3
51 – 60 (112 – 133)		40	0.4
61 – 75 (134 – 166)	50 mcg per 0.5 mL	50	0.5
76 – 85 (167 – 188)	80 mcg per 0.5 mL	64	0.4
>85 (>188)		80	0.5

* Must use vial. Minimum delivery for REDIPEN 0.3 mL

** When reconstituted as directed

2.4 Discontinuation of Dosing

Adults

It is recommended that patients receiving PegIntron, alone or in combination with ribavirin, be discontinued from therapy if HCV viral levels remain detectable after 6 months of therapy.

Pediatrics (3-17 years of age)

It is recommended that patients receiving PegIntron/REBETOL combination (excluding those with HCV Genotype 2 and 3) be discontinued from therapy at 12 weeks if their treatment week 12 HCV-RNA dropped $<2 \log_{10}$ compared to pretreatment or at 24 weeks if they have detectable HCV-RNA at treatment week 24.

2.5 Renal Function

In patients with moderate renal dysfunction (creatinine clearance 30-50 mL/min), the PegIntron dose should be reduced by 25%. Patients with severe renal dysfunction (creatinine clearance 10-29 mL/min) including those on hemodialysis, should have the PegIntron dose reduced by 50%. If renal function decreases during treatment, PegIntron therapy should be discontinued. When PegIntron is administered in combination with REBETOL, subjects with impaired renal function or those over the age of 50 should be more carefully monitored with respect to the development of anemia. PegIntron/REBETOL should not be used in patients with creatinine clearance <50 mL/min.

2.6 Preparation and Administration

PegIntron REDIPEN

PegIntron REDIPEN consists of a dual-chamber glass cartridge with sterile, lyophilized peginterferon alfa-2b in the active chamber and Sterile Water for Injection, USP in the diluent chamber. The PegIntron in the glass cartridge should appear as a white to off-white tablet shaped solid that is whole or in pieces, or powder.

To reconstitute the lyophilized peginterferon alfa-2b in the REDIPEN:

- Hold the REDIPEN upright (dose button down) and press the two halves of the pen together until there is an audible click.
- Gently invert the pen to mix the solution. **DO NOT SHAKE**. The reconstituted solution has a concentration of either 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL or 150 mcg per 0.5 mL for a single subcutaneous injection.
- Visually inspect the solution for particulate matter and discoloration prior to administration. The reconstituted solution should be clear and colorless. Do not use the solution if it is discolored, or not clear, or if particulates are present.

Keeping the pen upright, attach the supplied needle and select the appropriate PegIntron dose by pulling back on the dosing button until the dark bands are visible and turning the button until the dark band is aligned with the correct dose. The prepared PegIntron solution is to be injected subcutaneously.

The PegIntron REDIPEN is a single-use pen and does not contain a preservative. The reconstituted solution should be used immediately and cannot be stored for more than 24 hours at 2° - 8° C [see How Supplied/Storage and Handling (16)]. **DO NOT REUSE THE REDIPEN**. The sterility of any remaining product can no longer be guaranteed. **DISCARD THE UNUSED PORTION**. Pooling of unused portions of some medications has been linked to bacterial contamination and morbidity.

PegIntron Vials

Two B-D® Safety Lok™ syringes are provided in the package; one syringe is for the reconstitution steps and one for the patient injection. There is a plastic safety sleeve to be pulled over the needle after use. The syringe locks with an audible click when the green stripe on the safety sleeve covers the red stripe on the needle. Instructions for the preparation and administration of PegIntron Powder for Injection are provided below.

- **Reconstitute the PegIntron lyophilized product with only 0.7 mL of the 1.25 mL of supplied diluent (Sterile Water for Injection, USP). The diluent vial is for single use only. The remaining diluent should be discarded.** No other medications should be added to solutions containing PegIntron, and PegIntron should not be reconstituted with other diluents.
- Swirl gently to hasten complete dissolution of the powder. The reconstituted solution should be clear and colorless.
- Visually inspect the solution for particulate matter and discoloration prior to administration. The solution should not be used if discolored or cloudy, or if particulates are present.
- The appropriate PegIntron dose should be withdrawn and injected subcutaneously. PegIntron vials are for single use only and do not contain a preservative.

The reconstituted solution should be used immediately and cannot be stored for more than 24 hours at 2° - 8° C [see How Supplied/Storage and Handling (16)]. **DO NOT REUSE THE VIAL**. The sterility of any remaining product can no longer be guaranteed. **DISCARD THE UNUSED PORTION**. Pooling of unused portions of some medications has been linked to bacterial contamination and morbidity.

3 DOSAGE FORMS AND STRENGTHS

- Single-use vial: 1.25 mL diluent vial: 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL, 150 mcg per 0.5 mL
- Single-use REDIPEN: 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL, 150 mcg per 0.5 mL

4 CONTRAINDICATIONS

PegIntron is contraindicated in patients with:

- Known hypersensitivity reactions, such as urticaria, angioedema, bronchoconstriction, anaphylaxis, Stevens Johnson syndrome, and toxic epidermal necrolysis to interferon alpha or any other component of the product
- autoimmune hepatitis
- hepatic decompensation (Child-Pugh score >6 [class B and C]) in cirrhotic CHC patients before or during treatment.

PegIntron /REBETOL combination therapy is additionally contraindicated in:

- women who are pregnant. REBETOL may cause fetal harm when administered to a pregnant woman. REBETOL is contraindicated in women who are or may become pregnant. If this drug is used during pregnancy, or if the patient becomes pregnant while taking this drug the patient should be apprised of the potential hazard to a fetus. [see *Use in Specific Populations (8.1)*]
- men whose female partners are pregnant
- patients with hemoglobinopathies (e.g., thalassemia major, sickle-cell anemia)
- patients with creatinine clearance <50 mL/min

5 WARNINGS AND PRECAUTIONS

Patients should be monitored for the following serious conditions, some of which may become life threatening. Patients with persistently severe or worsening signs or symptoms should be withdrawn from therapy.

5.1 Use with Ribavirin

Pregnancy

REBETOL may cause birth defects and death of the unborn child. REBETOL therapy should not be started until a report of a negative pregnancy test has been obtained immediately prior to planned initiation of therapy. Patients should use at least two forms of contraception and have monthly pregnancy tests [see BOXED WARNING, Contraindications (4), Patient Counseling Information (17) and REBETOL package insert].

Anemia

Ribavirin caused hemolytic anemia in 10% of PegIntron/REBETOL- treated subjects within 1-4 weeks of initiation of therapy. Complete blood counts should be obtained pretreatment and at week 2 and week 4 of therapy or more frequently if clinically indicated. Anemia associated with REBETOL therapy may result in a worsening of cardiac disease. Decrease in dosage or discontinuation of REBETOL may be necessary [see *Dosage and Administration (2.3)* and *REBETOL package insert*].

5.2 Neuropsychiatric Events

Life-threatening or fatal neuropsychiatric events, including suicide, suicidal and homicidal ideation, depression, relapse of drug addiction/overdose, and aggressive behavior sometimes directed towards others have occurred in patients with and without a previous psychiatric disorder during PegIntron treatment and follow-up. Psychoses, hallucinations, bipolar disorders, and mania have been observed in patients treated with interferon alpha. PegIntron should be used with extreme caution in patients with a history of psychiatric disorders. Patients should be advised to report immediately any symptoms of depression or suicidal ideation to their prescribing physicians. Physicians should monitor all patients for evidence of depression and other psychiatric symptoms. If patients develop psychiatric problems, including clinical depression, it is recommended that the patients be carefully monitored during treatment and in the 6-month follow-up period. If psychiatric symptoms persist or worsen, or suicidal ideation or aggressive behavior towards others is identified, it is recommended that treatment with PegIntron be discontinued, and the patient followed, with psychiatric intervention as appropriate. In severe cases, PegIntron should be stopped immediately and psychiatric intervention instituted [see *Dosage and Administration (2.3)*]. Cases of encephalopathy have been observed in some patients, usually elderly, treated at higher doses of PegIntron.

5.3 Cardiovascular Events

Cardiovascular events, which include hypotension, arrhythmia, tachycardia, cardiomyopathy, angina pectoris, and myocardial infarction, have been observed in patients treated with PegIntron. PegIntron should be used cautiously in patients with cardiovascular disease. Patients with a history of myocardial infarction and arrhythmic disorder who require PegIntron therapy should be closely monitored [see *Warning and Precautions (5.15)*]. Patients with a history of significant or unstable cardiac disease should not be treated with PegIntron /REBETOL combination therapy [see *REBETOL package insert*].

5.4 Endocrine Disorders

PegIntron causes or aggravates hypothyroidism and hyperthyroidism. Hyperglycemia has been observed in patients treated with PegIntron. Diabetes mellitus has been observed in patients treated with alpha interferons. Patients with these conditions who cannot be effectively treated by medication should not begin PegIntron therapy. Patients who develop these conditions during treatment and cannot be controlled with medication should not continue PegIntron therapy.

5.5 Ophthalmologic Disorders

Decrease or loss of vision, retinopathy including macular edema, retinal artery or vein thrombosis, retinal hemorrhages and cotton wool spots, optic neuritis, and papilledema may be induced or aggravated by treatment with peginterferon alfa-2b or other alpha interferons. All patients should receive an eye examination at baseline. Patients with preexisting ophthalmologic disorders (e.g., diabetic or hypertensive retinopathy) should receive periodic ophthalmologic exams during interferon alpha treatment. Any patient who develops ocular symptoms should receive a prompt and complete eye examination. Peginterferon alfa-2b treatment should be discontinued in patients who develop new or worsening ophthalmologic disorders.

5.6 Cerebrovascular disorders

Ischemic and hemorrhagic cerebrovascular events have been observed in patients treated with interferon alfa-based therapies, including PegIntron. Events occurred in patients with few or no reported risk factors for stroke, including patients less than 45 years of age. Because these are spontaneous reports, estimates of frequency cannot be made and a causal relationship between interferon alfa based therapies and these events is difficult to establish.

5.7 Bone Marrow Toxicity

PegIntron suppresses bone marrow function, sometimes resulting in severe cytopenias. PegIntron should be discontinued in patients who develop severe decreases in neutrophil or platelet counts [see *Dosage and Administration (2.3)*]. Ribavirin may potentiate the neutropenia induced by interferon alpha. Very rarely alpha interferons may be associated with aplastic anemia.

5.8 Autoimmune Disorders

Development or exacerbation of autoimmune disorders (e.g., thyroiditis, thrombotic thrombocytopenic purpura, idiopathic thrombocytopenic purpura, rheumatoid arthritis, interstitial nephritis, systemic lupus erythematosus, and psoriasis) have been observed in patients receiving PegIntron. PegIntron should be used with caution in patients with autoimmune disorders.

5.9 Pancreatitis

Fatal and nonfatal pancreatitis have been observed in patients treated with alpha interferon. PegIntron therapy should be suspended in patients with signs and symptoms suggestive of pancreatitis and discontinued in patients diagnosed with pancreatitis.

5.10 Colitis

Fatal and nonfatal ulcerative or hemorrhagic/ischemic colitis have been observed within 12 weeks of the start of alpha interferon treatment. Abdominal pain, bloody diarrhea, and fever are the typical manifestations. PegIntron treatment should be discontinued immediately in patients who develop these signs and symptoms. The colitis usually resolves within 1-3 weeks of discontinuation of alpha interferons.

5.11 Pulmonary Disorders

Dyspnea, pulmonary infiltrates, pneumonia, bronchiolitis obliterans, interstitial pneumonitis, and sarcoidosis, some resulting in respiratory failure or patient deaths, may be induced or aggravated by PegIntron or alpha interferon therapy. Recurrence of respiratory failure has been observed with interferon rechallenge. PegIntron combination treatment should be suspended in patients who develop pulmonary infiltrates or pulmonary function impairment. Patients who resume interferon treatment should be closely monitored.

5.12 Hepatic Failure

Chronic hepatitis C (CHC) patients with cirrhosis may be at risk of hepatic decompensation and death when treated with alpha interferons, including PegIntron. Cirrhotic CHC patients coinfecting with HIV receiving highly active antiretroviral therapy (HAART) and alpha interferons with or without ribavirin appear to be at increased risk for the development of hepatic decompensation compared to patients not receiving HAART. During treatment, patients' clinical status and hepatic function should be closely monitored, and PegIntron treatment should be immediately discontinued if decompensation (Child-Pugh score >6) is observed [see *Contraindications (4)*].

5.13 Patients with Renal Insufficiency

Increases in serum creatinine levels have been observed in patients with renal insufficiency receiving interferon alpha products, including PegIntron. Patients with impaired renal function should be closely monitored for signs and symptoms of interferon toxicity, including increases in serum creatinine, and PegIntron dosing should be adjusted accordingly or discontinued [see *Clinical Pharmacology (12.3)* and *Dosage and Administration (2.3)*]. PegIntron monotherapy should be used with caution in patients with creatinine clearance <50 mL/min; the potential risks should be weighed against the potential benefits in these patients. Combination therapy with REBETOL must not be used in patients with creatinine clearance <50 mL/min [see *REBETOL Package Insert*].

5.14 Hypersensitivity

Serious, acute hypersensitivity reactions (e.g., urticaria, angioedema, bronchoconstriction, anaphylaxis) and cutaneous eruptions (Stevens Johnson syndrome, toxic epidermal necrolysis) have been rarely observed during alpha interferon therapy. If such a reaction develops during treatment with PegIntron, discontinue treatment and institute appropriate medical therapy immediately. Transient rashes do not necessitate interruption of treatment.

5.15 Laboratory Tests

PegIntron alone or in combination with ribavirin may cause severe decreases in neutrophil and platelet counts, and hematologic, endocrine (e.g., TSH), and hepatic abnormalities. Transient elevations in ALT (2- to 5-fold above baseline) were observed in 10% of subjects treated with PegIntron, and was not associated with deterioration of other liver functions. Triglyceride levels are frequently elevated in patients receiving alpha interferon therapy including PegIntron and should be periodically monitored.

Patients on PegIntron or PegIntron/REBETOL combination therapy should have hematology and blood chemistry testing before the start of treatment and then periodically thereafter. In the adult clinical trial CBC (including hemoglobin, neutrophil, and platelet counts) and chemistries (including AST, ALT, bilirubin, and uric acid) were measured during the treatment period at weeks 2, 4, 8, 12, and then at 6-week intervals or more frequently if abnormalities developed. In pediatric subjects the same laboratory parameters were evaluated with additional assessment of hemoglobin at treatment week 6. TSH levels were measured every 12 weeks during the treatment period. HCV RNA should be measured periodically during treatment [see *Dosage and Administration (2)*].

Patients who have pre-existing cardiac abnormalities should have electrocardiograms done before treatment with PegIntron/REBETOL.

5.16 Dental and Periodontal Disorders

Dental and periodontal disorders have been reported in patients receiving PegIntron/REBETOL combination therapy. In addition, dry mouth could have a damaging effect on teeth and mucous membranes of the mouth during long-term treatment with the combination of REBETOL and PegIntron. Patients should brush their teeth thoroughly twice daily and have regular dental examinations. If vomiting occurs, patients should be advised to rinse out their mouth thoroughly afterwards.

5.17 Triglycerides

Elevated triglyceride levels have been observed in patients treated with interferon alpha including PegIntron therapy. Hypertriglyceridemia may result in pancreatitis [see *Warnings and Precautions (5.9)*]. Elevated triglyceride levels should be managed as clinically appropriate. Discontinuation of PegIntron therapy should be considered for patients with symptoms of potential pancreatitis, such as abdominal pain, nausea, or vomiting and persistently elevated triglycerides (e.g., triglycerides >1000 mg/dL).

5.18 Impact on Growth-Pediatric Use

Data on the effects of PegIntron plus REBETOL on growth come from an open-label study in subjects 3 through 17 years of age, and weight and height changes are compared to U.S. normative population data. In general, the weight and height gain of pediatric subjects treated with PegIntron plus REBETOL lags behind that predicted by normative population data for the entire length of treatment. After about 6 months post-treatment (Follow-Up Week 24), subjects had weight gain rebounds and regained their weight to 53rd percentile, above the average of the normative population and similar to that predicted by their average baseline weight (57th percentile). After about 6 months post-treatment, height gain stabilized and subjects treated with PegIntron plus REBETOL had an average height percentile of 44th percentile, which was less than the average of the normative population and less than their average baseline height (51st percentile). Severely inhibited growth velocity (< 3rd percentile) was observed in 70% of the subjects while on treatment. Of the subjects experiencing severely inhibited growth, 20% had continued inhibited growth velocity (< 3rd percentile) after 6 months of follow-up.

Among the boys studied, the age groups of 3-11 years old and 12-17 years old had similar height percentile decreases of approximately 5 percentiles after 6 months post-treatment; weight gain continued to be similar to their average baseline percentile. Girls who were 3-11 years old and treated for 48 weeks had the largest average drop in height and weight percentiles (13 percentiles and 7 percentiles, respectively), whereas girls 12-17 years old continued along their average baseline height and weight percentiles after 6 months post-treatment.

6 ADVERSE REACTIONS

Clinical trials with PegIntron alone or in combination with REBETOL have been conducted in over 6900 subjects from 3 to 75 years of age

Serious adverse reactions have occurred in approximately 12% of subjects in clinical trials with PegIntron with or without REBETOL. [See **BOXED WARNING, Warnings and Precautions (5)**]. The most common serious events occurring in subjects treated with PegIntron and REBETOL were depression and suicidal ideation [see **Warnings and Precautions (5.2)**] each occurring at a frequency of less than 1%. The most common fatal events occurring in subjects treated with PegIntron and REBETOL were cardiac arrest, suicidal ideation and suicide attempt [see **Warnings and Precautions (5.2 and 5.5)**], all occurring in less than 1% of subjects.

Greater than 96% of all subjects in clinical trials experienced one or more adverse events. The most commonly reported adverse reactions in adult subjects receiving either PegIntron or PegIntron/REBETOL were injection site inflammation/reaction, fatigue/asthenia, headache, rigors, fevers, nausea, myalgia, and emotional lability/irritability. The most common adverse events in pediatric subjects, ages 3 and older, were pyrexia, headache, vomiting, neutropenia, fatigue, anorexia, injection site erythema, and abdominal pain.

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Adults

Study 1 compared PegIntron monotherapy with INTRON A monotherapy. Study 2 compared combination therapy of PegIntron/REBETOL with combination therapy with INTRON A/REBETOL. In these studies, nearly all study subjects in clinical trials experienced one or more adverse reactions.

Adverse reactions that occurred in Studies 1 and 2 at >5% incidence are provided in Table 8 by treatment group. Due to potential differences in ascertainment procedures, adverse reaction rate comparisons across studies should not be made.

Table 8
Adverse Reactions Occurring in > 5% of Subjects

Adverse Reactions	Percentage of Subjects Reporting Adverse Reactions*			
	Study 1		Study 2	
	PegIntron 1 mcg/kg (n=297)	INTRON A 3 MIU (n=303)	PegIntron 1.5 mcg/kg/ REBETOL (n=511)	INTRON A/ REBETOL (n=505)
Application Site				
Injection Site Inflammation/Reaction	47	20	75	49
Autonomic Nervous System				
Dry Mouth	6	7	12	8
Increased Sweating	6	7	11	7
Flushing	6	3	4	3
Body as a Whole				
Fatigue/Asthenia	52	54	66	63
Headache	56	52	62	58
Rigors	23	19	48	41
Fever	22	12	46	33
Weight Loss	11	13	29	20
Right Upper Quadrant Pain	8	8	12	6
Chest Pain	6	4	8	7
Malaise	7	6	4	6
Central/Peripheral Nervous System				
Dizziness	12	10	21	17
Endocrine				
Hypothyroidism	5	3	5	4
Gastrointestinal				
Nausea	26	20	43	33
Anorexia	20	17	32	27

Adverse Reactions	Percentage of Subjects Reporting Adverse Reactions*			
	Study 1		Study 2	
	PegIntron 1 mcg/kg (n=297)	INTRON A 3 MIU (n=303)	PegIntron 1.5 mcg/kg/ REBETOL (n=511)	INTRON A/ REBETOL (n=505)
Diarrhea	18	16	22	17
Vomiting	7	6	14	12
Abdominal Pain	15	11	13	13
Dyspepsia	6	7	9	8
Constipation	1	3	5	5
Hematologic Disorders				
Neutropenia	6	2	26	14
Anemia	0	0	12	17
Leukopenia	<1	0	6	5
Thrombocytopenia	7	<1	5	2
Liver and Biliary System				
Hepatomegaly	6	5	4	4
Musculoskeletal				
Myalgia	54	53	56	50
Arthralgia	23	27	34	28
Musculoskeletal Pain	28	22	21	19
Psychiatric				
Insomnia	23	23	40	41
Depression	29	25	31	34
Anxiety/Emotional Lability/Irritability	28	34	47	47
Concentration Impaired	10	8	17	21
Agitation	2	2	8	5
Nervousness	4	3	6	6
Reproductive, Female				
Menstrual Disorder	4	3	7	6
Resistance Mechanism				
Viral Infection	11	10	12	12
Fungal Infection	<1	3	6	1
Respiratory System				
Dyspnea	4	2	26	24
Coughing	8	5	23	16
Pharyngitis	10	7	12	13
Rhinitis	2	2	8	6
Sinusitis	7	7	6	5
Skin and Appendages				
Alopecia	22	22	36	32
Pruritus	12	8	29	28
Rash	6	7	24	23
Skin Dry	11	9	24	23
Special Senses, Other				
Taste Perversion	<1	2	9	4
Vision Disorders				
Vision Blurred	2	3	5	6
Conjunctivitis	4	2	4	5

*Subjects reporting one or more adverse reactions. A subject may have reported more than one adverse reaction within a body system/organ class category.

The adverse reaction profile in Study 3, which compared PegIntron/weight-based REBETOL combination to a PegIntron/flat dose REBETOL regimen, revealed an increased rate of anemia with weight-based dosing (29% vs. 19% for weight-based vs. flat dose regimens, respectively). However, the majority of cases of anemia were mild and responded to dose reductions.

The incidence of serious adverse reactions was comparable in all studies. In the PEG monotherapy trial (Study 1) the incidence of serious adverse reactions was similar (about 12%) in all treatment groups. In Study 2, the incidence of serious adverse reactions was 17% in the PegIntron/REBETOL groups compared to 14% in the INTRON A/REBETOL group. In Study 3, there was a similar incidence of serious adverse reactions reported for the weight-based REBETOL group (12%) and with the flat dose REBETOL regimen.

In many but not all cases, adverse reactions resolved after dose reduction or discontinuation of therapy. Some subjects experienced ongoing or new serious adverse reactions during the 6-month follow-up period. There have been 19 subject deaths which occurred during treatment or during follow-up in these clinical trials. In Study 1, there was one suicide in a subject receiving PegIntron monotherapy and two deaths among subjects receiving INTRON A monotherapy (1 murder/suicide and 1 sudden death). In Study 2, there was one suicide in a subject receiving PegIntron/REBETOL combination therapy; and 1 subject death in the INTRON A/REBETOL group (motor vehicle accident). In Study 3, there were 14 deaths, 2 of which were probable suicides and one was an unexplained death in a person with a relevant medical history of depression.

In Studies 1 and 2, 10-14% of subjects receiving PegIntron, alone or in combination with REBETOL, discontinued therapy compared with 6% treated with INTRON A alone and 13% treated with INTRON A in combination with REBETOL. Similarly in Study 3, 15% of subjects receiving PegIntron in combination with

weight-based REBETOL and 14% of subjects receiving PegIntron and flat dose REBETOL discontinued therapy due to an adverse reaction. The most common reasons for discontinuation of therapy were related to known interferon effects of psychiatric, systemic (e.g., fatigue, headache), or gastrointestinal adverse reactions.

In Study 2, dose reductions due to adverse reactions occurred in 42% of subjects receiving PegIntron (1.5 mcg/kg)/REBETOL and in 34% of those receiving INTRON A/REBETOL. The majority of subjects (57%) weighing 60 kg or less receiving PegIntron (1.5 mcg/kg)/REBETOL required dose reduction. Reduction of interferon was dose related (PegIntron 1.5 mcg/kg > PegIntron 0.5 mcg/kg or INTRON A), 40%, 27%, 28%, respectively. Dose reduction for REBETOL was similar across all three groups, 33-35%. The most common reasons for dose modifications were neutropenia (18%), or anemia (9%). Other common reasons included depression, fatigue, nausea, and thrombocytopenia. In Study 3, dose modifications due to adverse reactions occurred more frequently with WBD compared to flat dosing (29 and 23%, respectively).

In the PegIntron/REBETOL combination trials the most common adverse reactions were psychiatric which occurred among 77% of subjects in Study 2 and 68-69% of subjects in Study 3. These psychiatric adverse reactions included most commonly depression, irritability, and insomnia, each reported by approximately 30-40% of subjects in all treatment groups. Suicidal behavior (ideation, attempts, and suicides) occurred in 2% of all subjects during treatment or during follow-up after treatment cessation [see Warnings and Precautions (5.2)].

PegIntron induced fatigue or headache in approximately two-thirds of subjects, with fever or rigors in approximately half of the subjects. The severity of some of these systemic symptoms (e.g., fever and headache) tends to decrease as treatment continues. In Studies 1 and 2, application site inflammation and reaction (e.g., bruise, itchiness, and irritation) occurred at approximately twice the incidence with PegIntron therapies (in up to 75% of subjects) compared with INTRON A. However, injection site pain was infrequent (2-3%) in all groups. In Study 3 there was a 23-24% incidence overall for injection site reactions or inflammation.

In Study 2, many subjects continued to experience adverse reactions several months after discontinuation of therapy. By the end of the 6-month follow-up period, the incidence of ongoing adverse reactions by body class in the PegIntron 1.5/REBETOL group was 33% (psychiatric), 20% (musculoskeletal), and 10% (for endocrine and for GI). In approximately 10-15% of subjects weight loss, fatigue, and headache had not resolved.

Individual serious adverse reactions occurred at a frequency ≤1% and included suicide attempt, suicidal ideation, severe depression; psychosis, aggressive reaction, relapse of drug addiction/overdose; nerve palsy (facial, oculomotor); cardiomyopathy, myocardial infarction, angina, pericardial effusion, retinal ischemia, retinal artery or vein thrombosis, blindness, decreased visual acuity, optic neuritis, transient ischemic attack, supraventricular arrhythmias, loss of consciousness; neutropenia, infection (sepsis, pneumonia, abscess, cellulitis); emphysema, bronchiolitis obliterans, pleural effusion, gastroenteritis, pancreatitis, gout, hyperglycemia, hyperthyroidism and hypothyroidism, autoimmune thrombocytopenia with or without purpura, rheumatoid arthritis, interstitial nephritis, lupus-like syndrome, sarcoidosis, aggravated psoriasis; urticaria, injection-site necrosis, vasculitis, and phototoxicity.

Pediatric Subjects

In general, the adverse-reaction profile in the pediatric population was similar to that observed in adults. In the pediatric study, the most prevalent adverse reactions in all subjects were pyrexia (80%), headache (62%), neutropenia (33%), fatigue (30%), anorexia (29%), injection-site erythema (29%) and vomiting (27%). The majority of adverse reactions reported in the study were mild or moderate in severity. Severe adverse reactions were reported in 7% (8/107) of all subjects and included injection site pain (1%), pain in extremity (1%), headache (1%), neutropenia (1%), and pyrexia (4%). Important adverse reactions that occurred in this subject population were nervousness (7%; 7/107), aggression (3%; 3/107), anger (2%; 2/107), and depression (1%; 1/107). Five subjects received levothyroxine treatment, 3 with clinical hypothyroidism and 2 with asymptomatic TSH elevations.

Dose modifications were required in 25% of subjects, most commonly for anemia, neutropenia and weight loss. Two subjects (2%; 2/107) discontinued therapy as the result of an adverse reaction.

Adverse reactions that occurred with a ≥10% incidence in the pediatric trial subjects are provided in Table 9.

Table 9

Percentage of Pediatric Subjects With Treatment-Emergent Treatment-Related Adverse Reactions (in at Least 10% of All Subjects)

System Organ Class Preferred Term	All subjects n=107
Blood and Lymphatic System Disorders	
Neutropenia	33%
Anemia	11%
Leukopenia	10%
Gastrointestinal Disorders	
Abdominal Pain	21%
Abdominal Pain Upper	12%
Vomiting	27%
Nausea	18%
General Disorders and Administration Site Conditions	
Pyrexia	80%
Fatigue	30%
Injection Site Erythema	29%
Chills	21%
Asthenia	15%
Irritability	14%
Investigations	
Weight Decreased	19%
Metabolism and Nutrition Disorders	
Anorexia	29%
Decreased Appetite	22%
Musculoskeletal and Connective Tissue Disorders	
Arthralgia	17%
Myalgia	17%

Nervous System Disorders	
Headache	62%
Dizziness	14%
Skin and Subcutaneous Tissue Disorders	
Alopecia	17%

Laboratory Values

Adults

Changes in selected laboratory values during treatment with PegIntron alone or in combination with REBETOL treatment are described below. **Decreases in hemoglobin, neutrophils, and platelets may require dose reduction or permanent discontinuation from therapy** [See *Dosage and Administration (2.3)* and *Warnings and Precautions (5.1 and 5.7)*].

Hemoglobin. Hemoglobin levels decreased to <11 g/dL in about 30% of subjects in Study 2. In Study 3, 47% of subjects receiving WBD REBETOL and 33% on flat dose REBETOL had decreases in hemoglobin levels <11 g/dL. Reductions in hemoglobin to <9 g/dL occurred more frequently in subjects receiving WBD compared to flat dosing (4% and 2%, respectively). In study 2, dose modification was required in 9% and 13% of subjects in the PegIntron/REBETOL and INTRON A/REBETOL groups. Hemoglobin levels become stable by treatment week 4-6 on average. The typical pattern observed was a decrease in hemoglobin levels by treatment week 4 followed by stabilization and a plateau, which was maintained to the end of treatment. In the PegIntron monotherapy trial, hemoglobin decreases were generally mild and dose modifications were rarely necessary [See *Dosage and Administration (2.3)*].

Neutrophils. Decreases in neutrophil counts were observed in a majority of subjects treated with PegIntron alone (70%) or as combination therapy with REBETOL in Study 2 (85%) and INTRON A/REBETOL (60%). Severe potentially life-threatening neutropenia (<0.5 x 10⁹/L) occurred in 1% of subjects treated with PegIntron monotherapy, 2% of subjects treated with INTRON A/REBETOL, and in approximately 4% of subjects treated with PegIntron/REBETOL in Study 2. Two percent of subjects receiving PegIntron monotherapy and 18% of subjects receiving PegIntron/REBETOL in Study 2 required modification of interferon dosage. Few subjects (<1%) required permanent discontinuation of treatment. Neutrophil counts generally return to pretreatment levels 4 weeks after cessation of therapy [See *Dosage and Administration (2.3)*].

Platelets. Platelet counts decreased to <100,000/mm³ in approximately 20% of subjects treated with PegIntron alone or with REBETOL and in 6% of subjects treated with INTRON A/REBETOL. Severe decreases in platelet counts (<50,000/mm³) occur in <4% of subjects. Patients may require discontinuation or dose modification as a result of platelet decreases [See *Dosage and Administration (2.3)*]. In Study 2, 1% or 3% of subjects required dose modification of INTRON A or PegIntron, respectively. Platelet counts generally returned to pretreatment levels 4 weeks after the cessation of therapy.

Triglycerides. Elevated triglyceride levels have been observed in patients treated with interferon alphas including PegIntron [see *Warnings and Precautions (5.17)*].

Thyroid Function. Development of TSH abnormalities, with and without clinical manifestations, are associated with interferon therapies. In study 2, clinically apparent thyroid disorders occur among subjects treated with either INTRON A or PegIntron (with or without REBETOL) at a similar incidence (5% for hypothyroidism and 3% for hyperthyroidism). Subjects developed new onset TSH abnormalities while on treatment and during the follow-up period. At the end of the follow-up period, 7% of subjects still had abnormal TSH values [see *Warnings and Precautions (5.4)*].

Bilirubin and Uric Acid. In Study 2, 10-14% of subjects developed hyperbilirubinemia and 33-38% developed hyperuricemia in association with hemolysis. Six subjects developed mild to moderate gout.

Pediatric Subjects

Decreases in hemoglobin, white blood cells, platelets and neutrophils may require dose reduction or permanent discontinuation from therapy [See *Dosage and Administration (2.3)*]. Changes in selected laboratory values during treatment of 107 pediatric subjects with PegIntron/REBETOL combination therapy are described in Table 10. Most of the changes in laboratory values in this study were mild or moderate.

Table 10. Selected Hematological Abnormalities During Treatment Phase with PegIntron Plus REBETOL in Previously Untreated Pediatric Subjects

Laboratory Parameter ^a	All subjects n=107
Hemoglobin (g/dL)	
9.5 – <11.0	30%
8.0 – <9.5	2%
WBC (x10⁹/L)	
2.0 – 2.9	39%
1.5 – <2.0	3%
Platelets (x10⁹/L)	
70 – 100	1%
50 – <70	-
25 – <50	1%
Neutrophils (x10⁹/L)	
1.0 – 1.5	35%
0.75 – <1.0	26%
0.5 – <0.75	13%
<0.5	3%
Total Bilirubin	
1.26 - 2.59 xN ^b	7%
Evidence of Hepatic Failure	-

a: The table summarizes the worst category observed within the period per subject per laboratory test. Only subjects with at least one treatment value for a given laboratory test are included.

6.2 Immunogenicity

As with all therapeutic proteins, there is potential for immunogenicity. Approximately 2% of subjects receiving PegIntron (32/1759) or INTRON A (11/728) with or without REBETOL developed low-titer (≤ 160) neutralizing antibodies to PegIntron or INTRON A. The clinical and pathological significance of the appearance of serum neutralizing antibodies is unknown. The incidence of antibody formation is highly dependent on the sensitivity and specificity of the assay. Additionally, the observed incidence of antibody (including neutralizing antibody) positivity in an assay may be influenced by several factors including assay methodology, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to PegIntron with the incidence of antibodies to other products may be misleading.

6.3 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of PegIntron therapy: thrombotic thrombocytopenic purpura, pure red cell aplasia, erythema multiforme, Stevens Johnson syndrome, toxic epidermal necrolysis, seizures, memory loss, migraine headache, peripheral neuropathy, hearing impairment, hearing loss, vertigo, aphthous stomatitis, rhabdomyolysis, myositis, renal insufficiency, and renal failure. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

7 DRUG INTERACTIONS

7.1 Drugs Metabolized by Cytochrome P-450

When administering PegIntron with medications metabolized by CYP2C8/9 (e.g., warfarin and phenytoin) or CYP2D6 (e.g., flecainide), the therapeutic effect of these substrates may be decreased. [See *Clinical Pharmacology (12.3)*].

7.2 Methadone

PegIntron may increase methadone concentrations [see *Clinical Pharmacology (12.3)*]. The clinical significance of this finding is unknown; however, patients should be monitored for the signs and symptoms of increased narcotic effect.

7.3 Use with Ribavirin (Nucleoside Analogues)

Hepatic decompensation (some fatal) has occurred in cirrhotic HIV/HCV co-infected patients receiving combination antiretroviral therapy for HIV and interferon alpha and ribavirin. Adding treatment with alpha interferons alone or in combination with ribavirin may increase the risk in this patient subset. Patients receiving interferon with ribavirin and Nucleoside Reverse Transcriptase Inhibitors (NRTIs) should be closely monitored for treatment-associated toxicities, especially hepatic decompensation and anemia. Discontinuation of NRTIs should be considered as medically appropriate [see *Individual NRTI Product Information*]. Dose reduction or discontinuation of interferon, ribavirin, or both should also be considered if worsening clinical toxicities are observed, including hepatic decompensation (e.g., Child-Pugh >6).

Stavudine, Lamivudine, and Zidovudine

In vitro studies have shown ribavirin can reduce the phosphorylation of pyrimidine nucleoside analogues such as stavudine, lamivudine and zidovudine. In a study with another pegylated interferon alpha, no evidence of a pharmacokinetic or pharmacodynamic (e.g., loss of HIV/HCV virologic suppression) interaction was seen when ribavirin was co-administered with zidovudine, lamivudine or stavudine in HIV/HCV coinfecting subjects [See *Clinical Pharmacology (12.3)*].

HIV/HCV co-infected subjects who were administered zidovudine in combination with pegylated interferon alpha and ribavirin developed severe neutropenia (ANC <500) and severe anemia (hemoglobin <8 g/dL) more frequently than similar subjects not receiving zidovudine.

Didanosine

Co-administration of REBETOL Capsules or Oral Solution and didanosine is not recommended. Reports of fatal hepatic failure, as well as peripheral neuropathy, pancreatitis, and symptomatic hyperlactatemia/lactic acidosis have been reported in clinical trials [See *Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

PegIntron Monotherapy

Pregnancy Category C: Non-pegylated interferon alfa-2b, has been shown to have abortifacient effects in *Macaca mulatta* (rhesus monkeys) at 15 and 30 million IU/kg (estimated human equivalent of 5 and 10 million IU/kg, based on body surface area adjustment for a 60 kg adult). PegIntron should be assumed to also have abortifacient potential. *There* are no adequate and well-controlled studies in pregnant women. PegIntron therapy is to be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. Therefore, PegIntron is recommended for use in fertile women only when they are using effective contraception during the treatment period.

Use with Ribavirin

Pregnancy Category X: Significant teratogenic and/or embryocidal effects have been demonstrated in all animal species exposed to ribavirin. REBETOL therapy is contraindicated in women who are pregnant and in the male partners of women who are pregnant [See *Contraindications (4) and the REBETOL Package Insert*].

A Ribavirin Pregnancy Registry has been established to monitor maternal-fetal outcomes of pregnancies in female patients and female partners of male patients exposed to ribavirin during treatment and for 6 months following cessation of treatment. Physicians and patients are encouraged to report such cases by calling 1-800-593-2214.

8.3 Nursing Mothers

It is not known whether the components of PegIntron and/or REBETOL are excreted in human milk. Studies in mice have shown that mouse interferons are excreted in breast milk. Because of the potential for adverse reactions from the drug in nursing infants, a decision must be made whether to discontinue nursing or discontinue the PegIntron and REBETOL treatment, taking into account the importance of the therapy to the mother.

8.4 Pediatric Use

Safety and effectiveness in pediatric patients below the age of 3 years have not been established. Clinical trials in pediatric patients < 3 years of age are not considered feasible due to the small proportion of patients in this age group requiring treatment for CHC.

8.5 Geriatric Use

In general, younger patients tend to respond better than older patients to interferon-based therapies. Clinical studies of PegIntron alone or in combination with REBETOL did not include sufficient numbers of subjects aged 65 and over, however, to determine whether they respond differently than younger subjects. Treatment with alpha interferons, including PegIntron, is associated with neuropsychiatric, cardiac, pulmonary, GI, and systemic (flu-like) adverse effects. Because these adverse

reactions may be more severe in the elderly, caution should be exercised in the use of PegIntron in this population. This drug is known to be substantially excreted by the kidney. Because elderly patients are more likely to have decreased renal function, the risk of toxic reactions to this drug may be greater in patients with impaired renal function [see *Clinical Pharmacology (12.3)*]. When using PegIntron/REBETOL therapy, refer also to the REBETOL Package Insert.

8.6 Organ Transplant Recipients

The safety and efficacy of PegIntron alone or in combination with REBETOL for the treatment of hepatitis C in liver or other organ transplant recipients have not been studied. In a small (n=16) single-center, uncontrolled case experience, renal failure in renal allograft recipients receiving interferon alpha and ribavirin combination therapy was more frequent than expected from the center's previous experience with renal allograft recipients not receiving combination therapy. The relationship of the renal failure to renal allograft rejection is not clear.

8.7 HIV or HBV Coinfection

The safety and efficacy of PegIntron/REBETOL for the treatment of patients with HCV co-infected with HIV or HBV have not been established.

10 OVERDOSAGE

There is limited experience with overdosage. In the clinical studies, a few subjects accidentally received a dose greater than that prescribed. There were no instances in which a participant in the monotherapy or combination therapy trials received more than 10.5 times the intended dose of PegIntron. The maximum dose received by any subject was 3.45 mcg/kg weekly over a period of approximately 12 weeks. The maximum known overdosage of REBETOL was an intentional ingestion of 10 g (fifty 200 mg capsules). There were no serious reactions attributed to these overdosages. In cases of overdosing, symptomatic treatment and close observation of the patient are recommended.

11 DESCRIPTION

PegIntron, peginterferon alfa-2b, Powder for Injection is a covalent conjugate of recombinant alfa-2b interferon with monomethoxy polyethylene glycol (PEG). The average molecular weight of the PEG portion of the molecule is 12,000 daltons. The average molecular weight of the PegIntron molecule is approximately 31,000 daltons. The specific activity of peginterferon alfa-2b is approximately 0.7×10^8 IU/mg protein.

Interferon alfa-2b, is a water-soluble protein with a molecular weight of 19,271 daltons produced by recombinant DNA techniques. It is obtained from the bacterial fermentation of a strain of *Escherichia coli* bearing a genetically engineered plasmid containing an interferon gene from human leukocytes.

PegIntron is supplied in both vials and the REDIPEN for subcutaneous use.

Vials

Each vial contains either 74 mcg, 118.4 mcg, 177.6 mcg, or 222 mcg of PegIntron as a white to off-white tablet-like solid, that is whole/in pieces or as a loose powder, and 1.11 mg dibasic sodium phosphate anhydrous, 1.11 mg monobasic sodium phosphate dihydrate, 59.2 mg sucrose, and 0.074 mg polysorbate 80. Following reconstitution with 0.7 mL of the supplied Sterile Water for Injection, USP, each vial contains PegIntron at strengths of either 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL, or 150 mcg per 0.5 mL.

REDIPEN

REDIPEN is a dual-chamber glass cartridge containing lyophilized PegIntron as a white to off-white tablet or powder that is whole or in pieces in the sterile active chamber and a second chamber containing Sterile Water for Injection, USP. Each PegIntron REDIPEN contains either 67.5 mcg, 108 mcg, 162 mcg, or 202.5 mcg of PegIntron, and 1.013 mg dibasic sodium phosphate anhydrous, 1.013 mg monobasic sodium phosphate dihydrate, 54 mg sucrose, and 0.0675 mg polysorbate 80. Each cartridge is reconstituted to allow for the administration of up to 0.5 mL of solution. Following reconstitution, each REDIPEN contains PegIntron at strengths of either 50 mcg per 0.5 mL, 80 mcg per 0.5 mL, 120 mcg per 0.5 mL, or 150 mcg per 0.5 mL for a single use. Because a small volume of reconstituted solution is lost during preparation of PegIntron, each REDIPEN contains an excess amount of PegIntron powder and diluent to ensure delivery of the labeled dose.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Pegylated recombinant human interferon alfa-2b is an inducer of the innate antiviral immune response. [See *Clinical Pharmacology (12.4)*].

12.2 Pharmacodynamics

The pharmacodynamic effects of peginterferon alfa-2b include inhibition of viral replication in virus-infected cells, the suppression of cell cycle progression/cell proliferation, induction of apoptosis, anti-angiogenic activities, and numerous immunomodulating activities, such as enhancement of the phagocytic activity of macrophages, activation of NK cells, stimulation of cytotoxic T-lymphocytes, and the upregulation of the Th1 T-helper cell subset.

PegIntron raises concentrations of effector proteins such as serum neopterin and 2'5' oligoadenylate synthetase, raises body temperature, and causes reversible decreases in leukocyte and platelet counts. The correlation between the *in vitro* and *in vivo* pharmacologic and pharmacodynamic and clinical effects is unknown.

12.3 Pharmacokinetics

Following a single subcutaneous dose of PegIntron, the mean absorption half-life ($t_{1/2k_a}$) was 4.6 hours. Maximal serum concentrations (C_{max}) occur between 15-44 hours postdose, and are sustained for up to 48-72 hours. The C_{max} and AUC measurements of PegIntron increase in a dose-related manner. After multiple dosing, there is an increase in bioavailability of PegIntron. Week 48 mean trough concentrations (320 pg/mL; range 0, 2960) are approximately 3-fold higher than Week 4 mean trough concentrations (94 pg/mL; range 0, 416). The mean PegIntron elimination half-life is approximately 40 hours (range 22 to 60 hours) in patients with HCV infection. The apparent clearance of PegIntron is estimated to be approximately 22 mL/hr·kg. Renal elimination accounts for 30% of the clearance.

Pegylation of interferon alfa-2b produces a product (PegIntron) whose clearance is lower than that of non-pegylated interferon alfa-2b. When compared to INTRON A, PegIntron (1 mcg/kg) has approximately a 7-fold lower mean apparent clearance and a 5-fold greater mean half-life permitting a reduced dosing frequency. At effective therapeutic doses, PegIntron has approximately 10-fold greater C_{max} and 50-fold greater AUC than interferon alfa-2b.

Renal Dysfunction

Following multiple dosing of PegIntron (1 mcg/kg subcutaneously given every week for four weeks) the clearance of PegIntron is reduced by a mean of 17% in subjects with moderate renal impairment (creatinine clearance 30-49 mL/min) and by a mean of 44% in subjects with severe renal impairment (creatinine clearance 10-29 mL/min) compared to subjects with normal renal function. Clearance was similar in subjects with severe renal impairment not on dialysis and subjects who are receiving hemodialysis. The dose of PegIntron for monotherapy should be reduced in patients with moderate or severe renal impairment [see *Dosage and Administration (2.3)* and *REBETOL Package Insert*]. REBETOL should not be used in patients with creatinine clearance <50 mL/min [see *REBETOL Package Insert, WARNINGS*].

Gender

During the 48-week treatment period with PegIntron, no differences in the pharmacokinetic profiles were observed between male and female subjects with chronic hepatitis C infection.

Geriatric Patients

The pharmacokinetics of geriatric subjects (>65 years of age) treated with a single subcutaneous dose of 1 mcg/kg of PegIntron were similar in C_{max} , AUC, clearance, or elimination half-life as compared to younger subjects (28 to 44 years of age).

Pediatric Patients

Population pharmacokinetics for PegIntron and REBETOL (Capsules and Oral Solution) were evaluated in pediatric subjects with chronic hepatitis C between 3 and 17 years of age. In pediatric patients receiving PegIntron 60 mcg/m²/week subcutaneously, exposure may be approximately 50% higher than observed in adults receiving 1.5 mcg/kg/week subcutaneously. The pharmacokinetics of REBETOL (dose-normalized) in this trial were similar to those reported in a prior study of REBETOL in combination with INTRON A in pediatric subjects and in adults subjects.

Effect of Food on Absorption of Ribavirin

Both $AUC_{0-\infty}$ and C_{max} increased by 70% when REBETOL Capsules were administered with a high-fat meal (841 kcal, 53.8 g fat, 31.6 g protein, and 57.4 g carbohydrate) in a single-dose pharmacokinetic study [see *Dosage and Administration (2.2)*].

Drug Interactions

Drugs Metabolized by Cytochrome P-450

The pharmacokinetics of representative drugs metabolized by CYP1A2 (caffeine), CYP2C8/9 (tolbutamide), CYP2D6 (dextromethorphan), CYP3A4 (midazolam), and N-acetyltransferase (dapson) were studied in 22 subjects with chronic hepatitis C who received PegIntron (1.5 mcg/kg) once weekly for 4 weeks. PegIntron treatment resulted in a 28% (mean) increase in a measure of CYP2C8/9 activity. PegIntron treatment also resulted in a 66% (mean) increase in a measure of CYP2D6 activity; however, the effect was variable as 13 subjects had an increase, 5 subjects had a decrease, and 4 subjects had no significant change [see *Drug Interactions (7.1)*].

No significant effect was observed on the pharmacokinetics of representative drugs metabolized by CYP1A2, CYP3A4, or N-acetyltransferase. The effects of PegIntron on CYP2C19 activity were not assessed.

Methadone

The pharmacokinetics of concomitant administration of methadone and PegIntron were evaluated in 18 PegIntron naïve chronic hepatitis C subjects receiving 1.5 mcg/kg PegIntron subcutaneously weekly. All subjects were on stable methadone maintenance therapy receiving ≥ 40 mg/day prior to initiating PegIntron. Mean methadone AUC was approximately 16% higher after 4 weeks of PegIntron treatment as compared to baseline. In 2 subjects, methadone AUC was approximately double after 4 weeks of PegIntron treatment as compared to baseline [see *Drug Interactions (7.2)*].

Use with Ribavirin

Zidovudine, Lamivudine, and Stavudine

Ribavirin has been shown *in vitro* to inhibit phosphorylation of zidovudine, lamivudine, and stavudine. However, in a study with another pegylated interferon in combination with ribavirin, no pharmacokinetic (e.g., plasma concentrations or intracellular triphosphorylated active metabolite concentrations) or pharmacodynamic (e.g., loss of HIV/HCV virologic suppression) interaction was observed when ribavirin and lamivudine (n=18), stavudine (n=10), or zidovudine (n=6) were co-administered as part of a multi-drug regimen to HIV/HCV coinfecting subjects [see *Drug Interactions (7.3)*].

Didanosine

Exposure to didanosine or its active metabolite (dideoxyadenosine 5'-triphosphate) is increased when didanosine is co-administered with ribavirin, which could cause or worsen clinical toxicities [see *Drug Interactions (7.3)*].

12.4 Microbiology

Mechanism of Action

The biological activity of PegIntron is derived from its interferon alfa-2b moiety. Peginterferon alfa-2b binds to and activates the human type 1 interferon receptor. Upon binding, the receptor subunits dimerize, and activate multiple intracellular signal transduction pathways. Signal transduction is initially mediated by the JAK/STAT activation, which may occur in a wide variety of cells. Interferon receptor activation also activates NF κ B in many cell types. Given the diversity of cell types that respond to interferon alfa-2b, and the multiplicity of potential intracellular responses to interferon receptor activation, peginterferon alfa-2b is expected to have pleiotropic biological effects in the body.

The mechanism by which ribavirin contributes to its antiviral efficacy in the clinic is not fully understood. Ribavirin has direct antiviral activity in tissue culture against many RNA viruses. Ribavirin increases the mutation frequency in the genomes of several viruses and ribavirin triphosphate inhibits HCV polymerase in a biochemical reaction.

Antiviral Activity

The anti-HCV activity of interferon was demonstrated *in vitro* using self-replicating HCV RNA (HCV replicon cells) or HCV infection and resulted in an effective concentration (EC_{50}) value of 1-10 IU/mL.

The antiviral activity of ribavirin in the HCV-replicon is not well understood and has not been defined because of the cellular toxicity of ribavirin.

Resistance

HCV genotypes show wide variability in their response to pegylated recombinant human interferon/ribavirin therapy. Genetic changes associated with the variable response have not been identified.

Cross-resistance

There is no reported cross-resistance between pegylated/non-pegylated interferons and ribavirin.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis and Mutagenesis

PegIntron has not been tested for its carcinogenic potential. Neither PegIntron nor its components interferon or methoxypolyethylene glycol, caused damage to DNA when tested in the standard battery of mutagenesis assays, in the presence and absence of metabolic activation.

Use with Ribavirin: Ribavirin is genotoxic and mutagenic and should be considered a potential carcinogen. See REBETOL package insert for additional warnings relevant to PegIntron therapy in combination with ribavirin.

Impairment of Fertility

PegIntron may impair human fertility. Irregular menstrual cycles were observed in female cynomolgus monkeys given subcutaneous injections of 4239 mcg/m² PegIntron alone every other day for one month (approximately 345 times the recommended weekly human dose based upon body surface area). These effects included transiently decreased serum levels of estradiol and progesterone, suggestive of anovulation. Normal menstrual cycles and serum hormone levels resumed in these animals 2 to 3 months following cessation of PegIntron treatment. Every other day dosing with 262 mcg/m² (approximately 21 times the weekly human dose) had no effects on cycle duration or reproductive hormone status. The effects of PegIntron on male fertility have not been studied.

14 CLINICAL STUDIES

14.1 Chronic Hepatitis C in Adults

PegIntron Monotherapy-Study 1

A randomized study compared treatment with PegIntron (0.5, 1, or 1.5 mcg/kg once weekly subcutaneously) to treatment with INTRON A (3 million units three times weekly subcutaneously) in 1219 adults with chronic hepatitis from HCV infection. The subjects were not previously treated with interferon alpha, had compensated liver disease, detectable HCV RNA, elevated ALT, and liver histopathology consistent with chronic hepatitis. Subjects were treated for 48 weeks and were followed for 24 weeks posttreatment.

Seventy percent of all subjects were infected with HCV genotype 1, and 74 percent of all subjects had high baseline levels of HCV RNA (more than 2 million copies per mL of serum), two factors known to predict poor response to treatment.

Response to treatment was defined as undetectable HCV RNA and normalization of ALT at 24 weeks posttreatment. The response rates to the 1 and 1.5 mcg/kg PegIntron doses were similar (approximately 24%) to each other and were both higher than the response rate to INTRON A (12%). [See **Table 11**].

Table 11
Rates of Response to Treatment-Study 1

	A PegIntron 0.5 mcg/kg (N=315)	B PegIntron 1 mcg/kg (N=298)	C INTRON A 3 MIU three times weekly (N=307)	B - C (95% CI) Difference between PegIntron 1 mcg/kg and INTRON A
Treatment Response (Combined Virologic Response and ALT Normalization)	17%	24%	12%	11 (5, 18)
Virologic Response ^a	18%	25%	12%	12 (6,19)
ALT Normalization	24%	29%	18%	11 (5,18)

^a Serum HCV is measured by a research-based quantitative polymerase chain reaction assay by a central laboratory.

Subjects with both viral genotype 1 and high serum levels of HCV RNA at baseline were less likely to respond to treatment with PegIntron. Among subjects with the two unfavorable prognostic variables, 8% (12/157) responded to PegIntron treatment and 2% (4/169) responded to INTRON A. Doses of PegIntron higher than the recommended dose did not result in higher response rates in these subjects. Subjects receiving PegIntron with viral genotype 1 had a response rate of 14% (28/199) while subjects with other viral genotypes had a 45% (43/96) response rate.

Ninety-six percent of the responders in the PegIntron groups and 100% of responders in the INTRON A group first cleared their viral RNA by week 24 of treatment [see *Dosage and Administration (2)*].

The treatment response rates were similar in men and women. Response rates were lower in African American and Hispanic subjects and higher in Asians compared to Caucasians. Although African Americans had a higher proportion of poor prognostic factors compared to Caucasians, the number of non-Caucasians studied (9% of the total) was insufficient to allow meaningful conclusions about differences in response rates after adjusting for prognostic factors.

Liver biopsies were obtained before and after treatment in 60% of subjects. A modest reduction in inflammation compared to baseline that was similar in all four treatment groups was observed.

PegIntron/REBETOL Combination Therapy-Study 2

A randomized study compared treatment with two PegIntron/REBETOL regimens [PegIntron 1.5 mcg/kg subcutaneously once weekly/REBETOL 800 mg orally daily (in divided doses); PegIntron 1.5 mcg/kg subcutaneously once weekly for 4 weeks then 0.5 mcg/kg subcutaneously once weekly for 44 weeks/REBETOL 1000 or 1200 mg orally daily (in divided doses)] with INTRON A [3 MIU subcutaneously thrice weekly /REBETOL 1000/1200 mg orally daily (in divided doses)] in 1530 adults with chronic hepatitis C. Interferon-naïve subjects were treated for 48 weeks and followed for 24 weeks posttreatment. Eligible subjects had compensated liver disease, detectable HCV RNA, elevated ALT, and liver histopathology consistent with chronic hepatitis.

Response to treatment was defined as undetectable HCV RNA at 24 weeks posttreatment. The response rate to the PegIntron 1.5 mcg/kg plus ribavirin 800 mg dose was higher than the response rate to INTRON A/REBETOL [see **Table 12**]. The response rate to PegIntron 1.5→0.5 mcg/kg/REBETOL was essentially the same as the response to INTRON A/REBETOL (data not shown).

Table 12
Rates of Response to Treatment – Study 2

	PegIntron 1.5 mcg/kg once weekly REBETOL 800 mg daily	INTRON A 3 MIU three times weekly REBETOL 1000/1200 mg daily
Overall response ^{1,2}	52% (264/511)	46% (231/505)
Genotype 1	41% (141/348)	33% (112/343)
Genotype 2-6	75% (123/163)	73% (119/162)

¹ Serum HCV RNA is measured with a research-based quantitative polymerase chain reaction assay by a central laboratory.

² Difference in overall treatment response (PegIntron/REBETOL vs. INTRON A/REBETOL) is 6% with 95% confidence interval of (0.18, 11.63) adjusted for viral genotype and presence of cirrhosis at baseline. Response to treatment was defined as undetectable HCV RNA at 24 weeks posttreatment.

Subjects with viral genotype 1, regardless of viral load, had a lower response rate to PegIntron (1.5 mcg/kg)/REBETOL (800 mg) compared to subjects with other viral genotypes. Subjects with both poor prognostic factors (genotype 1 and high viral load) had a response rate of 30% (78/256) compared to a response rate of 29% (71/247) with INTRON A/REBETOL.

Subjects with lower body weight tended to have higher adverse reaction rates [see Adverse Reactions (6.1)] and higher response rates than subjects with higher body weights. Differences in response rates between treatment arms did not substantially vary with body weight.

Treatment response rates with PegIntron/REBETOL were 49% in men and 56% in women. Response rates were lower in African American and Hispanic subjects and higher in Asians compared to Caucasians. Although African Americans had a higher proportion of poor prognostic factors compared to Caucasians, the number of non-Caucasians studied (11% of the total) was insufficient to allow meaningful conclusions about differences in response rates after adjusting for prognostic factors in this study.

Liver biopsies were obtained before and after treatment in 68% of subjects. Compared to baseline, approximately 2/3 of subjects in all treatment groups were observed to have a modest reduction in inflammation.

PegIntron/REBETOL Combination Therapy-Study 3

In a large United States community-based study (Study 3), 4913 subjects with chronic hepatitis C were randomized to receive PegIntron 1.5 mcg/kg subcutaneously once weekly in combination with a REBETOL dose of 800-1400 mg (weight-based dosing- [WBD]) or 800 mg (Flat) orally daily (in divided doses) for 24 or 48 weeks based on genotype. Response to treatment was defined as undetectable HCV RNA (based on an assay with a lower limit of detection of 125 IU/mL) at 24 weeks posttreatment.

Treatment with PegIntron 1.5 mcg/kg and REBETOL 800-1400 mg resulted in a higher sustained virologic response compared to PegIntron in combination with a flat 800 mg daily dose of REBETOL. Subjects weighing >105 kg obtained the greatest benefit with WBD, although a modest benefit was also observed in subjects weighing >85-105 kg [see Table 13]. The benefit of WBD in subjects weighing >85 kg was observed with HCV genotypes 1-3. Insufficient data were available to reach conclusions regarding other genotypes. Use of WBD resulted in an increased incidence of anemia [see Adverse Reactions (6.1)].

Table 13
SVR Rate by Treatment and Baseline Weight- Study 3

Treatment Group	Subject Baseline Weight			
	<65 kg (<143 lb)	65-85 kg (143-188 lb)	>85-105 kg (>188-231 lb)	>105 kg (>231 lb)
WBD*	50% (173/348)	45% (449/994)	42% (351/835)	47% (138/292)
Flat	51% (173/342)	44% (443/1011)	39% (318/819)	33% (91/272)

* p=0.01, primary efficacy comparison (based on data from subjects weighing 65 kg or higher at baseline and utilizing a logistic regression analysis that includes treatment [WBD or Flat], genotype and presence/absence of advanced fibrosis, in the model).

A total of 1552 subjects weighing >65 kg in Study 3 had genotype 2 or 3 and were randomized to 24 or 48 weeks of therapy. No additional benefit was observed with the longer treatment duration.

14.2 Chronic Hepatitis C in Pediatrics

PegIntron/REBETOL Combination Therapy-Pediatric Study

Previously untreated pediatric subjects 3 to 17 years of age with compensated chronic hepatitis C and detectable HCV RNA were treated with REBETOL 15 mg/kg/day plus PegIntron 60 mcg/m² once weekly for 24 or 48 weeks based on HCV genotype and baseline viral load. All subjects were to be followed for 24 weeks post-treatment. A total of 107 subjects received treatment of whom 52% were female, 89% were Caucasian, and 67% were infected with HCV Genotype 1. Subjects infected with Genotype 1, 4 or Genotype 3 with HCV RNA ≥ 600,000 IU/mL received 48 weeks of therapy while those infected with Genotype 2 or Genotype 3 with HCV RNA < 600,000 IU/mL received 24 weeks of therapy. The study results are summarized in Table 14.

Table 14
Sustained Virologic Response Rates by Genotype and Treatment Duration – Pediatric Study

	All Subjects n=107	
	24 Weeks	48 Weeks
	Virologic Response n ^{a,b} (%)	Virologic Response n ^{a,b} (%)
Genotype		
All	26/27(96.3)	44/80(55.0)
1	-	38/72(52.8)
2	14/15(93.3)	-
3 ^c	12/12(100)	2/3(66.7)
4	-	4/5(80.0)

a: Response to treatment was defined as undetectable HCV RNA at 24 weeks posttreatment.

b: n = number of responders/number of subjects with given genotype, and assigned treatment duration.

c: Subjects with genotype 3 low viral load (<600,000 IU/mL) were to receive 24 weeks of treatment while those with genotype 3 and high viral load were to receive 48 weeks of treatment.

16 HOW SUPPLIED/STORAGE AND HANDLING

PegIntron REDIPEN

Each PegIntron REDIPEN Package Contains:	
A box containing one 50 mcg per 0.5 mL PegIntron REDIPEN and 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1323-01)
A box containing one 80 mcg per 0.5 mL PegIntron REDIPEN and 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1316-01)
A box containing one 120 mcg per 0.5 mL PegIntron REDIPEN and 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1297-01)
A box containing one 150 mcg per 0.5 mL PegIntron REDIPEN and 1B-D needle and 2 alcohol swabs.	(NDC 0085-1370-01)

Each PegIntron REDIPEN PAK 4 Contains:	
A box containing four 50 mcg per 0.5 mL PegIntron REDIPEN Units, each containing 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1323-02)
A box containing four 80 mcg per 0.5 mL PegIntron REDIPEN Units, each containing 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1316-02)
A box containing four 120 mcg per 0.5 mL PegIntron REDIPEN Units, each containing 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1297-02)
A box containing four 150 mcg per 0.5 mL PegIntron REDIPEN Units, each containing 1 B-D needle and 2 alcohol swabs.	(NDC 0085-1370-02)

PegIntron Vials

Each PegIntron Package Contains:	
A box containing one 50 mcg per 0.5 mL vial of PegIntron Powder for Injection and one 1.25 mL vial of Diluent (Sterile Water for Injection, USP), 2 B-D Safety Lok syringes with a safety sleeve and 2 alcohol swabs.	(NDC 0085-1368-01)
A box containing one 80 mcg per 0.5 mL vial of PegIntron Powder for Injection and one 1.25 mL vial of Diluent (Sterile Water for Injection, USP), 2 B-D Safety Lok syringes with a safety sleeve and 2 alcohol swabs.	(NDC 0085-1291-01)
A box containing one 120 mcg per 0.5 mL vial of PegIntron Powder for Injection and one 1.25 mL vial of Diluent (Sterile Water for Injection, USP), 2 B-D Safety Lok syringes with a safety sleeve and 2 alcohol swabs.	(NDC 0085-1304-01)
A box containing one 150 mcg per 0.5 mL vial of PegIntron Powder for Injection and one 1.25 mL vial of Diluent (Sterile Water for Injection, USP), 2 B-D Safety Lok syringes with a safety sleeve and 2 alcohol swabs.	(NDC 0085-1279-01)

Storage

PegIntron REDIPEN

PegIntron REDIPEN should be stored at 2°-8°C (36°-46°F).

After reconstitution, the solution should be used immediately, but may be stored up to 24 hours at 2°-8°C (36°-46°F). The reconstituted solution contains no preservative, and is clear and colorless. **DO NOT FREEZE.**

PegIntron Vials

PegIntron should be stored at 25°C (77°F); excursions permitted to 15°-30°C (59-86°F) [see USP Controlled Room Temperature]. After reconstitution with supplied Diluent the solution should be used immediately, but may be stored up to 24 hours at 2°-8°C (36°-46°F). The reconstituted solution contains no preservative, is clear and colorless. **DO NOT FREEZE.**

Disposal Instructions

Patients should be thoroughly instructed in the importance of proper disposal. After preparation and administration of the PegIntron for injection, patients should be advised to use a puncture-resistant container for the disposal of used syringes, needles, and the REDIPEN. The full container should be disposed of in accordance with state and local laws. Patients should also be cautioned against reusing or sharing needles, syringes, or the REDIPEN.

17 PATIENT COUNSELING INFORMATION

A patient should self-inject PegIntron only if it has been determined that it is appropriate and the patient agrees to medical follow-up as necessary and training in proper injection technique has been given to him/her.

17.1 Medication Guide

Patients receiving PegIntron alone or in combination with REBETOL should be directed in its appropriate use, informed of the benefits and risks associated with treatment, and referred to the MEDICATION GUIDES for PegIntron and, if applicable, REBETOL (ribavirin).

17.2 Pregnancy

Patients must be informed that REBETOL may cause birth defects and death of the unborn child. Extreme care must be taken to avoid pregnancy in female patients and in female partners of male patients during treatment with combination PegIntron/REBETOL therapy and for 6 months posttherapy. Combination PegIntron/REBETOL therapy should not be initiated until a report of a negative pregnancy test has been obtained immediately prior to initiation of therapy. It is recommended that patients undergo monthly pregnancy tests during therapy and for 6 months posttherapy [see *Contraindications (4)*, *Use in Specific Populations (8.1)* and *REBETOL package insert*].

17.3 HCV Transmission

Inform patients that there are no data regarding whether PegIntron therapy will prevent transmission of HCV infection to others. Also, it is not known if treatment with PegIntron will cure hepatitis C or prevent cirrhosis, liver failure, or liver cancer that may be the result of infection with the hepatitis C virus.

17.4 Laboratory Evaluations, Hydration, “Flu-like” Symptoms

Patients should be advised that laboratory evaluations are required before starting therapy and periodically thereafter [see *Warnings and Precautions (5.15)*]. It is advised that patients be well hydrated, especially during the initial stages of treatment. “Flu-like” symptoms associated with administration of PegIntron may be minimized by bedtime administration of PegIntron or by use of antipyretics.

Manufactured by Schering Corporation, a subsidiary of Schering-Plough Corporation, Kenilworth, NJ 07033 USA.



© 2003, 2005, 2006, Schering Corporation. All rights reserved.

*B-D and Safety-Lok are trademarks of Becton Dickinson and Company.

Rev 12/08
B-32063403
31852803T

1 **27853560T**

2 **MEDICATION GUIDE**

3 **PegIntron®**

4 **Peginterferon alfa-2b**

6 **Including appendix with instructions for using PegIntron® Powder for Injection**

8 Read this Medication Guide carefully before you start taking PegIntron (**Peg In-tron**) or
9 PegIntron/REBETOL (**REB-eh-tole**) combination therapy. Read the Medication Guide each
10 time you refill your prescription because there may be new information. The information in
11 this Medication Guide does not take the place of talking with your health care provider
12 (doctor, nurse, nurse practitioner, or physician's assistant).

14 **If you are taking PegIntron/REBETOL combination therapy, also read the Medication
15 Guide for REBETOL (ribavirin, USP) Capsules and Oral Solution.**

17 **What is the most important information I should know about PegIntron and
18 PegIntron/REBETOL combination therapy?**

20 PegIntron (peginterferon) is a treatment for some people who are infected with hepatitis C
21 virus. However, PegIntron and PegIntron/REBETOL combination therapy can have serious
22 side effects that may cause death in rare cases. Before you decide to start treatment, you
23 should talk to your health care provider about the possible benefits and side effects of
24 PegIntron or PegIntron/REBETOL combination therapy. If you begin treatment you will
25 need to see your health care provider regularly for medical examinations and lab tests to
26 make sure your treatment is working and to check for side effects.

28 **REBETOL may cause birth defects and/or death of an unborn child. If you are
29 pregnant, you or your male partner must not take PegIntron/REBETOL combination
30 therapy. You must not become pregnant while either you or your partner are being
31 treated with the combination PegIntron/REBETOL therapy, or for 6 months after
32 stopping therapy. Men and women should use birth control while taking the
33 combination therapy and for 6 months afterwards. If you or your partner are being
34 treated and you become pregnant either during treatment or within 6 months of
35 stopping treatment, call your health care provider right away. There is a Ribavirin
36 Pregnancy Registry that collects information about pregnancy outcomes in female
37 patients and female partners of male patients exposed to ribavirin. You or your
38 healthcare provider are encouraged to contact the Registry at 1-800-593-2214.**

40 **If you are taking PegIntron or PegIntron/REBETOL therapy you should call your
41 health care provider immediately if you develop any of these symptoms:**

43 **New or worsening mental health problems such as thoughts about killing or hurting
44 yourself or others, trouble breathing, chest pain, severe stomach or lower back pain,
45 bloody diarrhea or bloody bowel movements, high fever, bruising, bleeding, or
46 decreased vision.**



47
48 The most serious possible side effects of PegIntron and PegIntron/REBETOL therapy
49 include:

50

51 **Problems with Pregnancy.** Combination PegIntron/REBETOL therapy can cause
52 death, serious birth defects, or other harm to your unborn child. If you are a woman of
53 childbearing age you must not become pregnant during treatment and for 6 months
54 after you have stopped therapy. You must have a negative pregnancy test immediately
55 before beginning treatment, during treatment and for 6 months after you have stopped
56 therapy. Both male and female patients must use effective forms of birth control during
57 treatment and for the 6 months after treatment is completed. Male patients should use
58 a condom. If you are a female, you must use birth control even if you believe that you are
59 not fertile or that your fertility is low. You should talk to your health care provider about
60 birth control for you and your partner.

61

62 **Mental health problems and suicide.** PegIntron and PegIntron/REBETOL therapies may
63 cause patients to develop mood or behavioral problems. These can include irritability
64 (getting easily upset) and depression (feeling low, feeling bad about yourself, or feeling
65 hopeless). Some patients may have aggressive behavior. Former drug addicts may fall back
66 into drug addiction or overdose. Some patients think about hurting or killing themselves or
67 other people and some have killed (suicide) or hurt themselves or others. You must tell your
68 health care provider if you are being treated for a mental illness or had treatment in the past
69 for any mental illness, including depression and suicidal behavior. You should tell your
70 health care provider if you have ever been addicted to drugs or alcohol.

71

72 **Heart problems.** Some patients taking PegIntron or PegIntron/REBETOL therapy may
73 develop problems with their heart, including low blood pressure, fast heart rate, and very
74 rarely, heart attacks. Tell your health care provider if you have had any heart problems in the
75 past.

76

77 **Blood problems.** PegIntron and PegIntron/REBETOL therapies commonly lower two types
78 of blood cells (white blood cells and platelets). In some patients, these blood counts may fall
79 to dangerously low levels. If your blood counts become very low, this could lead to
80 infections or bleeding.

81

82 REBETOL therapy causes a decrease in the number of red blood cells you have (anemia).
83 This can be dangerous, especially for patients who already have heart or circulatory
84 (cardiovascular) problems. Talk with your health care provider before taking combination
85 PegIntron/REBETOL therapy if you have or have ever had any cardiovascular problems.

86

87 **Body organ problems.** Certain symptoms like severe stomach pain may mean that your
88 internal organs are being damaged. Cases of weakness, loss of coordination, and numbness
89 due to stroke have been reported in patients taking PegIntron, including patients with few or
90 no reported risk factors for stroke.

91

92

93 *For other possible side effects, see “What are the possible side effects of PegIntron and*
 94 *PegIntron/REBETOL” in this Medication Guide.*

95

96 **What is PegIntron and PegIntron/REBETOL combination therapy?**

97 The PegIntron product is a drug used to treat adults who have a lasting (chronic) infection
 98 with hepatitis C virus and who show signs that the virus is damaging the liver.

99 PegIntron/REBETOL combination therapy consists of two medications also used to treat
 100 hepatitis C infection in adults and children 3 years of age and older. Patients with hepatitis C
 101 have the virus in their blood and in their liver. PegIntron reduces the amount of virus in the
 102 body and helps the body's immune system fight the virus. REBETOL (ribavirin) is a drug
 103 that helps to fight the viral infection but does not work when used by itself to treat chronic
 104 hepatitis C.

105

106 It is not known if PegIntron or PegIntron/REBETOL therapies can cure hepatitis C
 107 (permanently eliminate the virus) or if it can prevent liver failure or liver cancer that is
 108 caused by hepatitis C infection.

109

110 It is also not known if PegIntron or PegIntron/REBETOL combination therapy will prevent
 111 one infected person from infecting another person with hepatitis C.

112

113 **Who should not take PegIntron or PegIntron/REBETOL therapy?**

114 Do not take PegIntron or PegIntron/REBETOL therapy if you:

115 • are pregnant, planning to get pregnant during treatment or during the 6 months after
 116 treatment, or breast-feeding

117

118 • are a male patient with a female sexual partner who is pregnant or plans to become
 119 pregnant at any time while you are being treated with REBETOL or during the 6
 120 months after your treatment has ended

121

122 • have hepatitis caused by your immune system attacking your liver (autoimmune
 123 hepatitis) or unstable liver disease

124

125 • had an allergic reaction to another alpha interferon or are allergic to any of the
 126 ingredients in PegIntron or REBETOL Capsules or Oral Solution. If you have any
 127 doubts, ask your health care provider

128

129 • Do not take PegIntron/REBETOL combination therapy if you have abnormal red
 130 blood cells such as is seen in sickle-cell anemia or thalassemia major

131

132 **If you have any of the following conditions or serious medical problems, discuss them**
 133 **with your health care provider before taking PegIntron or PegIntron/REBETOL**
 134 **therapy:**

135 • depression or anxiety

136 • sleep problems

137 • high blood pressure

138 • previous heart attack, or other heart problems

- 139 • liver problems (other than hepatitis C infection)
- 140 • any kind of autoimmune disease (where the body's immune system attacks the body's
- 141 own cells), such as psoriasis, systemic lupus erythematosus, rheumatoid arthritis
- 142 • thyroid problems
- 143 • diabetes
- 144 • colitis (inflammation of the bowels)
- 145 • cancer
- 146 • hepatitis B infection
- 147 • HIV infection
- 148 • kidney problems
- 149 • bleeding problems
- 150 • alcoholism
- 151 • drug abuse or addiction
- 152 • body organ transplant and are taking medicine that keeps your body from rejecting your
- 153 transplant (suppresses your immune system)

154

155

156 **How should I take PegIntron or PegIntron/REBETOL?**

157 Your health care provider will decide whether you will take PegIntron therapy alone or the
 158 combination of PegIntron/REBETOL, as well as the correct dose (for adults the dose of
 159 PegIntron is based on weight). For children 3 years of age and older, your healthcare
 160 provider will recommend the dose of PegIntron based on a calculation accounting for both
 161 height and weight. PegIntron and PegIntron/REBETOL are given for up to one year. Take
 162 your prescribed dose of PegIntron ONCE A WEEK, on the same day of each week and at
 163 approximately the same time. Take the medicine for the full course of prescribed therapy
 164 and do not take more than the prescribed dose. REBETOL should be taken with food.
 165 When you take REBETOL with food, more of the medicine (70% more on average) is taken
 166 up by your body. You should take REBETOL the same way every day (twice a day with
 167 food) to keep the medicine in your body at a steady level. This will help your health care
 168 provider to decide how your treatment is working and how to change the dose of REBETOL
 169 you take if you have side effects from REBETOL. **Be sure to read the Medication Guide**
 170 **for REBETOL (ribavirin, USP) for complete instructions on how to take the**
 171 **REBETOL capsules and oral solution.**

172

173 You should be completely comfortable with how to prepare PegIntron; how to set the dose
 174 you take, and how to inject yourself before you use PegIntron for the first time. PegIntron
 175 comes in two different forms, a powder in a single-use vial and a REDIPEN[®] single-use
 176 delivery system. See the attached appendix for detailed instructions for preparing and giving
 177 a dose of PegIntron.

178

179 If you miss a dose of the PegIntron product, take the missed dose as soon as possible during
 180 the same day or the next day, then continue on your regular dosing schedule. If several days
 181 go by after you miss a dose, check with your health care provider about what to do. Do not
 182 double the next dose or take more than one dose a week without talking to your health care
 183 provider. Call your health care provider right away if you take more than your prescribed

184 PegIntron dose. Your health care provider may wish to examine you more closely, and take
185 blood for testing.

186

187 If you miss a dose of REBETOL, take the missed dose as soon as possible during the same
188 day. If an entire day has gone by, check with your health care provider about what to do. Do
189 not double the next dose.

190

191 You must get regular blood tests to help your health care provider check how the treatment is
192 working and to check for side effects.

193

194 Tell your health care provider if you are taking or planning to take other prescription or non-
195 prescription medicines, including vitamin and mineral supplements and herbal medicines.

196

197 **What should I avoid while taking PegIntron or PegIntron/REBETOL therapies?**

198 • If you are pregnant do not start taking PegIntron/REBETOL combination therapy.

199 • Avoid becoming pregnant while taking PegIntron or PegIntron/REBETOL.

200 PegIntron and PegIntron/REBETOL may harm your unborn child (death or serious birth
201 defects) or cause you to lose your baby (miscarry). **If you or your partner become**
202 **pregnant during treatment or during the 6 months after treatment with**
203 **PegIntron/REBETOL combination therapy, immediately report the pregnancy to your**
204 **health care provider. You or your health care provider should call 1-800-593-2214.** By
205 calling this number, information about you and/or your partner will be added to a pregnancy
206 registry that will be used to help you and your health care provider make decisions about
207 your treatment for hepatitis in the future. You, your partner and/or your health care provider
208 will be asked to provide follow-up information on the outcome of the pregnancy.

209

- Do not breast-feed your baby while taking PegIntron.

210

211
212 **What are the possible side effects of PegIntron and PegIntron/REBETOL combination**
213 **therapy?**

214

215 **Possible, serious side effects include:**

216

- **Mental health problems including suicide, blood problems, heart problems, body organ problems.** See “What is the most important information I should know about PegIntron and PegIntron/REBETOL combination therapy?”

217

218

219

220

- **Other body organ problems.** A few patients have lung problems (such as pneumonia or inflammation of the lung tissue), inflammation of the kidney, and eye disorders.

221

222

223

224

- **New or worsening autoimmune disease.** Some patients taking PegIntron or PegIntron/REBETOL develop autoimmune diseases (a condition where the body’s immune cells attack other cells or organs in the body), including rheumatoid arthritis, systemic lupus erythematosus, and psoriasis. In some patients who already have an autoimmune disease, the disease worsens on PegIntron and PegIntron/REBETOL combination therapy.

225

226

227

228

229

230

231 **Common but less serious side effects include:**

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

- **Flu-like symptoms.** Most patients who take PegIntron or PegIntron/REBETOL therapy have "flu-like" symptoms (headache, muscle aches, tiredness and fever). Some of these symptoms (fever, headache) usually lessen after the first few weeks of therapy. You can reduce some of these symptoms by injecting your PegIntron dose at bedtime. Over-the-counter pain and fever reducers, such as acetaminophen or ibuprofen, can be used to prevent or reduce the fever and headache.
- **Extreme fatigue (tiredness).** Many patients become extremely tired while on PegIntron or PegIntron/REBETOL combination therapy.
- **Appetite problems.** Nausea, loss of appetite, and weight loss occur commonly.
- **Thyroid problems.** Some patients develop changes in the function of their thyroid. Symptoms of thyroid changes include the inability to concentrate, feeling cold or hot all the time, a change in your weight and changes to your skin.
- **Blood sugar problems.** Some patients develop problems with the way their body controls their blood sugar and may develop high blood sugar or diabetes.
- **Skin reactions.** Redness, swelling, and itching are common at the site of injection. If after several days these symptoms do not disappear contact your health care provider. You may get a rash during therapy. If this occurs, your health care provider may recommend medicine to treat the rash.
- **Hair thinning.** Hair thinning is common during PegIntron and PegIntron/REBETOL treatment. Hair loss stops and hair growth returns after therapy is stopped.
- **Effect on Growth in Children.** Weight loss and slowed growth are common in children during treatment with PegIntron/REBETOL. Catch-up weight gain and some catch-up in growth occur after the end of treatment, but some children may not reach their pre-treatment expected height.

These are not all of the side effects of PegIntron or PegIntron/REBETOL combination therapy. Your health care provider or pharmacist can give you a more complete list.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

General advice about prescription medicines:

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. If you have any concerns about PegIntron, ask your health care provider. Your health care provider or pharmacist can give you information about PegIntron that was written for



275 health care professionals. Do not use PegIntron for a condition for which it was not
276 prescribed. Do not share this medication with other people.

277

278 **If you are taking PegIntron/REBETOL combination therapy, also read the Medication**
279 **Guide for REBETOL (ribavirin, USP) Capsules and Oral Solution.**

280

281 *This Medication Guide has been approved by the U.S. Food and Drug Administration.*

282 **Revised December 2008**

283

284 **How do I prepare and inject the PegIntron Dose?**

285 Before you inject PegIntron, the powder must be mixed with **0.7 mL** of the supplied
286 DILUENT for PegIntron, Sterile Water for Injection (diluent). This product can also be
287 administered by a parent or care-taker as instructed by your healthcare provider. You should
288 carefully follow the directions given to you by your health care provider.

289

290 The vial of mixed PegIntron should be used immediately. **DO NOT** prepare more than one
291 vial at a time. If you don't use the vial of the prepared solution right away, it must be stored
292 in a refrigerator and used within 24 hours.

293

294 **Storing PegIntron**

295 PegIntron Powder should be stored at room temperature (25 ° C, 77°F); avoid exposure to
296 heat. After mixing, the PegIntron solution should be used immediately but may be stored in
297 the refrigerator up to 24 hours. The solution contains no preservatives. **DO NOT FREEZE.**

298 **Preparing the PegIntron solution:**

299

300 1. Find a clean, well-lit, non-slip flat working surface and assemble all of the supplies
301 you will need for an injection. All of the supplies you will need for an injection are
302 in the PegIntron Powder for Injection package. The package contains:

- 303 ▪ a vial of PegIntron powder
- 304
- 305 ▪ a 1.25 mL vial of DILUENT
- 306
- 307 ▪ 2 disposable syringes, and
- 308
- 309 ▪ alcohol swabs

310

311 2. Check the date printed on the PegIntron carton to make sure that the expiration date
312 has not passed. Remove one vial and look at the contents. The PegIntron in the vial
313 should appear as a white to off-white tablet-like solid that is whole/in pieces or as a
314 loose powder.

315

316 If you have already mixed the PegIntron solution and it has been stored properly in the
317 refrigerator, take it out of the refrigerator and allow the solution to come to room
318 temperature.

319



320 3. Wash your hands thoroughly with soap and water, rinse and towel dry. It is important
 321 to keep your work area, your hands and injection site clean to minimize the risk of
 322 infection.

323

324 The disposable syringes have needles that are already attached and cannot be removed.
 325 Each syringe has a clear plastic safety sleeve that is pulled over the needle for disposal
 326 after use. The safety sleeve should remain tight against the flange while using the
 327 syringe and moved over the needle only when ready for disposal. **Figure A.**

328 The syringes and needles are for single use only.

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346




Figure A

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

4. Remove the protective wrapper from ONE of the syringes provided and use for the following steps 5-7. Make sure that the syringe safety sleeve is sitting against the flange. (see **Figure A**).

5. Remove the protective plastic cap from the tops of both the supplied DILUENT and the PegIntron vials. Clean the rubber stopper on the top of both vials with an alcohol swab.

6. Carefully remove the protective cap straight off of the needle to avoid damaging the needle point. Fill the syringe with air by pulling the plunger to 0.7 mL (**Figure B**). Hold the DILUENT vial upright. Do not touch the cleaned top of the vial with your hands (**Figure C**). Insert the needle through the center of the rubber stopper of the DILUENT vial, and inject the air from the syringe into the vial (**Figure D**). Turn the vial upside down and make sure the tip of the needle is in the liquid. Withdraw only 0.7 mL of DILUENT by pulling the plunger back to the 0.7 mL mark on the side of the syringe (**Figure E**). Remove the needle from the vial (**Figure F**). **Discard the remaining DILUENT.**




Figure B




Figure C




Figure D




Figure E



Figure F

363
364365
366

367 7. Insert the needle through the center of the rubber stopper of the PegIntron vial, and place
368 the needle tip against the glass wall of the vial (**Figure G**). SLOWLY inject the 0.7 mL
369 DILUENT so that the stream of DILUENT runs down the side of the vial. To prevent
370 bubbles from forming, DO NOT AIM THE STREAM of diluent directly on the tablet-like
371 SOLID or POWDER in the bottom of the vial. Remove the needle from the vial.

372 Firmly grasp the safety sleeve and pull it over the exposed needle until you hear a click. The
373 green stripe on the safety sleeve will completely cover the red stripe on the needle. (See
374 **Figure O** in the section: "Injecting the PegIntron dose.") Discard the syringe and needle in
375 the puncture proof container.

376

- 377 8. GENTLY swirl the vial in a gentle circular motion (**Figure H**), until the PegIntron is
378 completely dissolved. **DO NOT SHAKE** the vial. If any powder remains undissolved in
379 the vial, gently turn the vial upside down until all of the powder is dissolved. It is not
380 unusual for the solution to appear cloudy or bubbly for a few minutes. If air bubbles do
381 form, wait until the solution has settled and all bubbles have risen to the top before
382 withdrawing your dose from the vial.

383

384

385

386

387

388

389

390

391

392

393

394






Figure G




Figure H

- 395 9. After the solution has settled and is completely dissolved it should be clear, colorless
 396 and without particles, but there may be a ring of foam or bubbles on the surface, this is
 397 normal. Do not use it if you see particles or the color is not correct.
- 398 10. After the PegIntron powder is dissolved but before you withdraw your dose, clean the
 399 rubber stopper again with an alcohol swab.
 400
 401
 402
 403
 404 .
- 405 11. Unwrap the second syringe provided. You will use it to give yourself the injection.
 406 Carefully remove the protective cap from the needle and fill the syringe with air by pulling
 407 the plunger to the number on the side of the syringe (mL) that corresponds to your prescribed
 408 dose (**Figure J**). Hold the PegIntron vial upright. DO NOT touch the cleaned top of the vial
 409 with your hands (**Figure K**). Insert the needle into the vial containing the PegIntron solution
 410 and inject the air into the center of the vial (**Figure L**).
 411


412
413
414 Figure J415
416 Figure K

417 Figure L

- 417 12. Turn the PegIntron vial upside down. Be sure the tip of the needle is in the PegIntron
 418 solution. While holding the vial and syringe with one hand slowly pull the plunger back to
 419 withdraw the exact amount of PegIntron into the syringe your health care provider told you
 420 to use (**Figure M**).
 421
 422

423
424 Figure M
425
426

427 13. Remove the needle from the vial (**Figure N**) and check for air bubbles in the syringe. If
 428 you see any bubbles, hold the syringe with the needle pointing up and gently tap the syringe
 429 until the bubbles rise. Then push the plunger in slowly until the bubbles disappear.
 430



431 Figure N

432

433

434 **Injecting the PegIntron Dose**

435 **Selecting the Site for Injection.**

436 The best sites for giving yourself an injection are those areas with a layer of fat between the
 437 skin and muscle, like your thigh, the outer surface of your upper arm, and abdomen. Do not
 438 inject yourself in the area near your navel or waistline. If you are very thin, you should only
 439 use the thigh or outer surface of the arm for injection.
 440

440

441 You should use a different site each time you inject PegIntron to avoid soreness at any one
 442 site. Do not inject PegIntron solution into an area where the skin is irritated, red, bruised,
 443 infected or has scars, stretch marks or lumps.
 444

444

445 14. Clean the skin where the injection is to be given with an alcohol swab, and wait for the
 446 area to dry. Remove the protective cap from the needle. Make sure the safety sleeve of the
 447 syringe is pushed firmly against the syringe flange so that the needle is fully exposed (see
 448 **Figure A**).
 449

449

450 15. With one hand, pinch a 2-inch fold of loose skin. With your other hand, pick up the
 451 syringe and hold it like a pencil. Position the bevel of the needle facing up and insert the
 452 needle approximately ¼ inch into the pinched skin at approximately a 45 to 90 degree angle
 453 with a quick dart-like thrust. After the needle is in, remove the hand that you used to pinch
 454 your skin and use it to hold the syringe barrel. Pull the plunger of the syringe back very
 455 slightly. If blood comes into the syringe, the needle has entered a blood vessel. **Do not**
 456 **inject.** Withdraw the needle and discard the syringe as outlined in step 17. Repeat the above
 457 steps with a new vial to prepare a new syringe and inject the medicine at a new site. If no
 458 blood is present in the syringe, inject the medicine by gently pressing the plunger all the way
 459 down the syringe barrel.
 460

460


461 16. Hold an alcohol swab near the needle and pull the needle straight out of the skin. Press
 462 the alcohol swab over the injection site for several seconds. Do not massage the injection
 463 site. If there is bleeding, cover it with a bandage.
 464

464

465 17. After injecting your dose, firmly grasp the safety sleeve and pull it over the exposed
 466 needle until you hear a click, and the green stripe on the safety sleeve covers the red stripe on

467 the needle (**Figure O**). Discard the syringe and needle in the Sharp's container supplied to
 468 you.

469
 470
 471
 472
 473
 474
 475
 476
 477



478 **Figure O**

479 18. After 2 hours, check the injection site for redness, swelling, or tenderness. If you have a
 480 skin reaction and it doesn't clear up in a few days, contact your health care provider or nurse.

481

482 **How do I dispose of the used syringes and needles?**

483 Discard used safety lock syringes and needles in a Sharp's container or other puncture proof
 484 container like a coffee can. **DO NOT USE** glass or clear plastic containers. Your health care
 485 provider or nurse will tell you how to dispose of a full container. Always keep the container
 486 out of reach of children.

487

488 Manufactured by: Schering Corporation, Kenilworth, NJ 07033 USA

489 **DATE 12/08**

490 **27853560T**

1 **27662455T**

2 **MEDICATION GUIDE**

3 **PegIntron™ REDIPEN® Single-dose Delivery System**
4 **(Peginterferon alfa-2b)**

5
6 **Including appendix with instructions for using PegIntron™ REDIPEN® Single-dose**
7 **Delivery System**

8
9 Read this Medication Guide carefully before you start taking PegIntron™ (**Peg In-tron**) or
10 PegIntron™/REBETOL® (**REB-eh-tole**) combination therapy. Read the Medication Guide
11 each time you refill your prescription because there may be new information. The
12 information in this Medication Guide does not take the place of talking with your health care
13 provider (doctor, nurse, nurse practitioner, or physician's assistant).

14
15 If you are taking PegIntron/REBETOL combination therapy, also read the Medication Guide
16 for REBETOL (ribavirin, USP) Capsules and Oral Solution.

17
18 **What is the most important information I should know about PegIntron and**
19 **PegIntron/REBETOL combination therapy?**

20
21 PegIntron (peginterferon) is a treatment for some people who are infected with hepatitis C
22 virus. However, PegIntron and PegIntron/REBETOL combination therapy can have serious
23 side effects that may cause death in rare cases. Before you decide to start treatment, you
24 should talk to your health care provider about the possible benefits and side effects of
25 PegIntron or PegIntron/REBETOL combination therapy. If you begin treatment you will
26 need to see your health care provider regularly for medical examinations and lab tests to
27 make sure your treatment is working and to check for side effects.

28
29 **REBETOL may cause birth defects and/or death of an unborn child. If you are**
30 **pregnant, you or your male partner must not take PegIntron/REBETOL combination**
31 **therapy. You must not become pregnant while either you or your partner are being**
32 **treated with the combination PegIntron/REBETOL therapy, or for 6 months after**
33 **stopping therapy. Men and women should use birth control while taking the**
34 **combination therapy and for 6 months afterwards. If you or your partner are being**
35 **treated and you become pregnant, either during treatment or within 6 months of**
36 **stopping treatment, call your health care provider right away. There is a Ribavirin**
37 **Pregnancy Registry that collects information about pregnancy outcomes of female**
38 **patients and female partners of male patients exposed to ribavirin. You or your**
39 **healthcare provider are encouraged to contact the Registry at 1-800-593-2214.**

40
41 **If you are taking PegIntron or PegIntron/REBETOL therapy you should call your**
42 **health care provider immediately if you develop any of these symptoms:**

43
44 **New or worsening mental health problems, such as thoughts about killing or hurting**
45 **yourself or others, trouble breathing, chest pain, severe stomach or lower back pain,**

46 **bloody diarrhea or bloody bowel movements, high fever, bruising, bleeding, or**
47 **decreased vision.**

48

49 The most serious possible side effects of PegIntron and PegIntron/REBETOL therapy
50 include:

51

52 **Problems with Pregnancy. Combination PegIntron/REBETOL therapy can cause**
53 **death, serious birth defects, or other harm to your unborn child. If you are a woman of**
54 **childbearing age, you must not become pregnant during treatment and for 6 months**
55 **after you have stopped therapy. You must have a negative pregnancy test immediately**
56 **before beginning treatment, during treatment, and for 6 months after you have stopped**
57 **therapy. Both males and female patients must use effective forms of birth control**
58 **during treatment and for the 6 months after treatment is completed. Male patients**
59 **should use a condom.** If you are a female, you must use birth control even if you believe
60 that you are not fertile or that your fertility is low. You should talk to your health care
61 provider about birth control for you and your partner.

62

63 **Mental health problems and suicide.** PegIntron and PegIntron/REBETOL therapies may
64 cause patients to develop mood or behavioral problems. These can include irritability (getting
65 easily upset) and depression (feeling low, feeling bad about yourself, or feeling hopeless).
66 Some patients may have aggressive behavior. Former drug addicts may fall back into drug
67 addiction or overdose. Some patients think about hurting or killing themselves or other
68 people and some have killed (suicide) or hurt themselves or others. You must tell your
69 health care provider if you are being treated for a mental illness or had treatment in the past
70 for any mental illness, including depression and suicidal behavior. You should tell your
71 health care provider if you have ever been addicted to drugs or alcohol.

72

73 **Heart problems.** Some patients taking PegIntron or PegIntron/REBETOL therapy may
74 develop problems with their heart, including low blood pressure, fast heart rate, and very
75 rarely, heart attacks. Tell your health care provider if you have had any heart problems in the
76 past.

77

78 **Blood problems.** PegIntron and PegIntron/REBETOL therapies commonly lower two types
79 of blood cells (white blood cells and platelets). In some patients, these blood counts may fall
80 to dangerously low levels. If your blood counts become very low, this could lead to
81 infections or bleeding.

82

83 REBETOL therapy causes a decrease in the number of red blood cells you have (anemia).
84 This can be dangerous, especially for patients who already have heart or circulatory
85 (cardiovascular) problems. Talk with your health care provider before taking combination
86 PegIntron/REBETOL therapy if you have, or have ever had any cardiovascular problems.

87

88 **Body organ problems.** Certain symptoms like severe stomach pain may mean that your
89 internal organs are being damaged. Cases of weakness, loss of coordination, and numbness
90 due to stroke have been reported in patients taking PegIntron, including patients with few or
91 no reported risk factors for stroke.

92
93 *For other possible side effects, see “What are the possible side effects of PegIntron and*
94 *PegIntron/REBETOL” in this Medication Guide.*
95

96 **What is PegIntron and PegIntron/REBETOL combination therapy?**

97 The PegIntron product is a drug used to treat adults who have a lasting (chronic) infection
98 with hepatitis C virus and who show signs that the virus is damaging the liver.
99 PegIntron/REBETOL combination therapy consists of two medications also used to treat
100 hepatitis C infection in adults and children 3 years of age and older. Patients with hepatitis C
101 have the virus in their blood and in their liver. PegIntron reduces the amount of virus in the
102 body and helps the body's immune system fight the virus. REBETOL (ribavirin) is a drug
103 that helps to fight the viral infection, but does not work when used by itself to treat chronic
104 hepatitis C.
105

106 It is not known if PegIntron or PegIntron/REBETOL therapies can cure hepatitis C
107 (permanently eliminate the virus), or if it can prevent liver failure or liver cancer that is
108 caused by hepatitis C infection.
109

110 It is also not known if PegIntron or PegIntron/REBETOL combination therapy will prevent
111 one infected person from infecting another person with hepatitis C.
112

113 **Who should not take PegIntron or PegIntron/REBETOL therapy?**

114 Do not take PegIntron or PegIntron/REBETOL therapy if you:

- 115 • are pregnant, planning to get pregnant during treatment or during the 6 months after
116 treatment, or breast-feeding
117
- 118 • are a male patient with a female sexual partner who is pregnant, or plans to become
119 pregnant at any time while you are being treated with REBETOL, or during the 6
120 months after your treatment has ended
121
- 122 • have hepatitis caused by your immune system attacking your liver (autoimmune
123 hepatitis) or unstable liver disease
124
- 125 • had an allergic reaction to another alpha interferon or are allergic to any of the
126 ingredients in PegIntron or REBETOL Capsules or Oral Solution. If you have any
127 doubts, ask your health care provider.
128
- 129 • Do not take PegIntron/REBETOL combination therapy if you have abnormal red
130 blood cells such as is seen in sickle-cell anemia or thalassemia major.
131

132 **If you have any of the following conditions or serious medical problems, discuss them**
133 **with your health care provider before taking PegIntron or PegIntron/REBETOL**
134 **therapy:**

- 135 • depression or anxiety
- 136 • sleep problems
- 137 • high blood pressure

-
- 138 • previous heart attack, or other heart problems
 - 139 • liver problems (other than hepatitis C infection)
 - 140 • any kind of autoimmune disease (where the body's immune system attacks the body's
 - 141 own cells), such as psoriasis, systemic lupus erythematosus, rheumatoid arthritis
 - 142 • thyroid problems
 - 143 • diabetes
 - 144 • colitis (inflammation of the bowels)
 - 145 • cancer
 - 146 • hepatitis B infection
 - 147 • HIV infection
 - 148 • kidney problems
 - 149 • bleeding problems
 - 150 • alcoholism
 - 151 • drug abuse or addiction
 - 152 • body organ transplant and are taking medicine that keeps your body from rejecting your
 - 153 transplant (suppresses your immune system)

154
155

156 **How should I take PegIntron or PegIntron/REBETOL?**

157 Your health care provider will decide whether you will take PegIntron therapy alone or the
158 combination of PegIntron/REBETOL, as well as the correct dose (for adults the dose of
159 PegIntron is based on weight). For children 3 years of age and older, your healthcare
160 provider will recommend the dose of PegIntron based on a calculation accounting for both
161 height and weight. PegIntron and PegIntron/REBETOL are given for up to one year. Take
162 your prescribed dose of PegIntron ONCE A WEEK, on the same day of each week and at
163 approximately the same time. Take the medicine for the full course of prescribed therapy and
164 do not take more than the prescribed dose. REBETOL should be taken with food. When you
165 take REBETOL with food, more of the medicine (70% more on average) is taken up by your
166 body. You should take REBETOL the same way every day (twice a day with food) to keep
167 the medicine in your body at a steady level. This will help your health care provider to decide
168 how your treatment is working and how to change the dose of REBETOL you take if you
169 have side effects from REBETOL. **Be sure to read the Medication Guide for REBETOL**
170 **(ribavirin, USP) for complete instructions on how to take the REBETOL capsules and**
171 **oral solution.**

172

173 You should be completely comfortable with how to prepare PegIntron, how to set the dose
174 you take, and how to inject yourself before you use PegIntron for the first time. PegIntron
175 comes in two different forms, a powder in a single-use vial and a REDIPEN single-use
176 delivery system. See the attached appendix for detailed instructions for preparing and giving
177 a dose of PegIntron.

178

179 If you miss a dose of the PegIntron product, take the missed dose as soon as possible during
180 the same day or the next day, then continue on your regular dosing schedule. If several days
181 go by after you miss a dose, check with your health care provider about what to do. Do not
182 double the next dose or take more than one dose a week without talking to your health care
183 provider. Call your health care provider right away if you take more than your prescribed

184 PegIntron dose. Your health care provider may wish to examine you more closely, and take
185 blood for testing.

186

187 If you miss a dose of REBETOL, take the missed dose as soon as possible during the same
188 day. If an entire day has gone by, check with your health care provider about what to do. Do
189 not double the next dose.

190

191 You must get regular blood tests to help your health care provider check how the treatment is
192 working and to check for side effects.

193

194 Tell your health care provider if you are taking or planning to take other prescription or non-
195 prescription medicines, including vitamin and mineral supplements and herbal medicines.

196

197 **What should I avoid while taking PegIntron or PegIntron/REBETOL therapies?**

198 • If you are pregnant do not start taking PegIntron/REBETOL combination therapy.

199 • Avoid becoming pregnant while taking PegIntron or PegIntron/REBETOL.

200 PegIntron and PegIntron/REBETOL may harm your unborn child (death or serious birth
201 defects) or cause you to lose your baby (miscarry). **If you or your partner becomes**
202 **pregnant during treatment or during the 6 months after treatment with**
203 **PegIntron/REBETOL combination therapy, immediately report the pregnancy to your**
204 **health care provider. You or your health care provider should call 1-800-593-2214.** By
205 calling this number, information about you and/or your partner will be added to a pregnancy
206 registry that will be used to help you and your health care provider make decisions about
207 your treatment for hepatitis in the future. You, your partner, and/or your health care provider
208 will be asked to provide follow-up information on the outcome of the pregnancy.

209

210 • Do not breast-feed your baby while taking PegIntron.

211

212 **What are the possible side effects of PegIntron and PegIntron/REBETOL combination**
213 **therapy?**

214

215 **Possible, serious side effects include:**

216 **Mental health problems including suicide, blood problems, heart problems, body organ**
217 **problems.** See “What is the most important information I should know about PegIntron and
218 PegIntron/REBETOL combination therapy?”

219

220 **Other body organ problems.** A few patients have lung problems (such as pneumonia or
221 inflammation of the lung tissue), inflammation of the kidney, and eye disorders.

222

223 **New or worsening autoimmune disease.** Some patients taking PegIntron or
224 PegIntron/REBETOL develop autoimmune diseases (a condition where the body’s immune
225 cells attack other cells or organs in the body), including rheumatoid arthritis, systemic lupus
226 erythematosus, and psoriasis. In some patients who already have an autoimmune disease, the
227 disease worsens on PegIntron and PegIntron/REBETOL combination therapy.

228

229 Common but less serious side effects include:

230

231 **Flu-like symptoms.** Most patients who take PegIntron or PegIntron/REBETOL therapy have
232 "flu-like" symptoms (headache, muscle aches, tiredness, and fever). Some of these symptoms
233 (fever, headache) usually lessen after the first few weeks of therapy. You can reduce some of
234 these symptoms by injecting your PegIntron dose at bedtime. Over-the-counter pain and
235 fever reducers, such as acetaminophen or ibuprofen, can be used to prevent or reduce the
236 fever and headache.

237

238 **Extreme fatigue (tiredness).** Many patients become extremely tired while on PegIntron or
239 PegIntron/REBETOL combination therapy.

240

241 **Appetite problems.** Nausea, loss of appetite, and weight loss, occur commonly.

242

243 **Thyroid problems.** Some patients develop changes in the function of their thyroid.
244 Symptoms of thyroid changes include the inability to concentrate, feeling cold or hot all the
245 time, a change in your weight, and changes to your skin.

246

247 **Blood sugar problems.** Some patients develop problems with the way their body controls
248 their blood sugar, and may develop high blood sugar or diabetes.

249

250 **Skin reactions.** Redness, swelling, and itching are common at the site of injection. If after
251 several days these symptoms do not disappear contact your health care provider. You may
252 get a rash during therapy. If this occurs, your health care provider may recommend medicine
253 to treat the rash.

254

255 **Hair thinning.** Hair thinning is common during PegIntron and PegIntron/REBETOL
256 treatment. Hair loss stops and hair growth returns after therapy is stopped.

257

258 **Effect on Growth in Children.** Weight loss and slowed growth are common in children
259 during treatment with PegIntron/REBETOL. Catch-up weight gain and some catch-up in
260 growth occur after the end of treatment, but some children may not reach their pre-treatment
261 expected height..

262

263 These are not all of the side effects of PegIntron or PegIntron/REBETOL combination
264 therapy. Your health care provider or pharmacist can give you a more complete list.

265

266 Call your doctor for medical advice about side effects. You may report side effects to FDA at
267 1-800-FDA-1088.

268

269 **General advice about prescription medicines:**

270 Medicines are sometimes prescribed for purposes other than those listed in a Medication
271 Guide. If you have any concerns about PegIntron, ask your health care provider. Your health
272 care provider or pharmacist can give you information about PegIntron that was written for
273 health care professionals. Do not use PegIntron for a condition for which it was not
274 prescribed. Do not share this medication with other people.

275

276 If you are taking PegIntron/REBETOL combination therapy, also read the Medication Guide
277 for REBETOL (ribavirin, USP) Capsules and Oral Solution.

278
279 *This Medication Guide has been approved by the U.S. Food and Drug Administration.*
280 Revised December 2008

281

282 **How do I prepare and inject the PegIntron REDIPEN Dose?**

283

284 The PegIntron REDIPEN system is for a single use, by one person only, ONCE A WEEK.
285 The REDIPEN must not be shared. Use only the injection needle provided in the packaging
286 for the PegIntron REDIPEN system. If you have problems with the REDIPEN system or the
287 PegIntron solution, you should contact your health care provider or pharmacist.

288

289 The following instructions explain how to prepare and inject yourself with the PegIntron
290 REDIPEN system. This product can also be administered by a parent or care-taker as
291 instructed by your healthcare provider. Please read the instructions carefully and follow them
292 step by step. Your health care provider will instruct you on how to self-inject with the
293 PegIntron REDIPEN . Do not attempt to inject yourself unless you are sure you understand
294 the procedure and requirements for self-injection.


295

296

297 **How to Use the PegIntron REDIPEN Single-dose Delivery System.**

298

299



300

301

302

303 **Storing PegIntron**

304 PegIntron REDIPEN should be stored in the refrigerator at 2°C to 8°C (36°F to 46°F); avoid
305 exposure to heat. After mixing, the PegIntron solution should be used immediately but may
306 be stored in the refrigerator up to 24 hours at 2°C to 8°C (36°F - 46°F). The solution contains
307 no preservatives. **DO NOT FREEZE.**

308

309 **Preparation**

310

311 1. Find a clean, well-lit, non-slip flat working surface and assemble all of the supplies you
312 will need for an injection. All of the supplies you will need are in the PegIntron
313 REDIPEN package. The package contains:

314

315

- a PegIntron REDIPEN single-dose delivery system
- one disposable needle

- 316
- 317
- 318
- two alcohol swabs, and
 - Dosing tray; (The dosing tray is the bottom half of the REDIPEN package.)
- 319
- 320
- 321
- 322
- 323
- 324
- 325
- 326
- 327
- 328
- 329
2. Take the PegIntron REDIPEN out of the refrigerator and allow the medicine to come to room temperature. Before removing the REDIPEN from the carton, check the expiration date printed on the PegIntron REDIPEN carton to make sure that the expiration date has not passed. Do not use if the expiration date has passed.
 3. After taking the PegIntron REDIPEN out of the carton, look in the window of the REDIPEN and make sure the PegIntron in the cartridge holder window is a white, to off white tablet that is whole, or in pieces, or powdered.
 4. Wash your hands thoroughly with soap and water, rinse, and towel dry. It is important to keep your work area, your hands, and the injection site clean to minimize the risk of infection.

330 1. Mix the Drug

331 **Key points:**

332 **Before you mix the PegIntron, make sure it is at room temperature. It is important that**

333 **you keep the PegIntron REDIPEN UPRIGHT (Dosing Button down) as shown in**

334 **Figure 1.**

337

338 a. Hold the PegIntron REDIPEN **UPRIGHT (Figure 1a)** in the dosing tray on a hard,

339 flat, non-slip surface with the dosing button **down**. You may want to hold the REDIPEN

340 using the grip.

341

342 b. To mix the powder and the liquid, keep the REDIPEN upright in the dosing tray and

343 press the top half of the REDIPEN downward toward the hard, flat, non-slip surface **until**

344 **you hear the click** (Figure 1b). Once you've heard the click, you will notice in the

345 window that both dark stoppers are now touching. The dosing button should be flush with


346 the pen body.



347 **Figure 1a**

348

349



350
351 **Figure 1b**

352
353 c. Wait several seconds for the powder to completely dissolve.

354
355
356 **d. Gently turn the PegIntron REDIPEN upside down twice (Figure 2). To avoid**
357 **excessive foaming, DO NOT SHAKE.**



358
359 **Figure 2**

360
361 e. Keep the PegIntron™ REDIPEN® **UPRIGHT**, with the dosing button down. Then,
362 look through the REDIPEN® window to see that the mixed PegIntron™ solution is
363 completely dissolved. The solution should be clear and colorless **before use**. Before
364 attaching the needle, it is normal to see some small bubbles in the REDIPEN® window,
365 near the top of the solution. Do not use the solution if it is discolored, or not clear, or if
366 particulates are present.

367
368 **f. Place the PegIntron REDIPEN back into the dosing tray provided in the**
369 **packaging (Figure 3). The dosing button will be on the bottom.**
370



Figure 3

371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388

2. Attach the Needle

- a. Wipe the rubber membrane of the PegIntron REDIPEN with one alcohol swab.
- b. Remove the protective paper tab from the injection needle, but do NOT remove either the outer cap or the yellow inner cap from the injection needle. Keeping the PegIntron REDIPEN UPRIGHT in the dosing tray, FIRMLY push the injection needle straight into the REDIPEN rubber membrane, and screw it firmly in place, in a clockwise direction (Figure 4). Remember to leave the needle caps in place when you attach the needle to the REDIPEN. Pushing the needle through the rubber membrane, “primes” the needle and allows the extra liquid and air in the pen to be removed.



389

Figure 4

390

391

392 NOTE: Some fluid will trickle out. This is **normal**. The dark stoppers move up and you will no
393 longer see the fluid in the window once the needle is successfully primed.

394

395 3. Dialing the Dose

396

397 a. **Remove the PegIntron REDIPEN from the dosing tray (Figure 5a).**

398 Holding the PegIntron REDIPEN firmly, pull the dosing button out as far as it will go.

399 You will see a dark band:

400

401 **Do not push the dosing button in until you are ready to self-inject the PegIntron**
402 **dose.**



403

404

405 **Figure 5a**


406

407 b. Turn the dosing button until your prescribed dose is lined up with the dosing tab (**Figure 5b**).

408 The dosing button will turn freely. If you have trouble dialing your dose, check to make sure the
409 dosing button has been pulled out **as far** as it will go (**Figure 5c**).

410

411



412

413

414 **Figure 5b**

Figure 5c

414

415 c. Carefully lay the PegIntron REDIPEN down on a hard, flat, non-slip surface. Do NOT
416 remove either of the needle caps and do NOT push the dosing button in until you are ready to
417 self-inject the PegIntron dose.
418

419 **4. Injecting the PegIntron Dose**

420 **Choosing an Injection Site**

421 The best sites for giving yourself an injection are those areas with a layer of fat between
422 the skin and muscle, like your thigh, the outer surface of your upper arm, and abdomen.
423 Do not inject yourself in the area near your navel or waistline. If you are very thin, you
424 should only use the thigh or outer surface of the arm for injection.
425

426 You should use a different site each time you inject PegIntron to avoid soreness at any
427 one site. Do not inject PegIntron into an area where the skin is irritated, red, bruised,
428 infected, or has scars, stretch marks, or lumps.
429

430 a. Clean the skin where the injection is to be given with the second alcohol swab provided,
431 and wait for the area to dry.

432 b. Remove the **outer** cap from the needle (**Figure 6a**). There may be some liquid around the
433 yellow inner needle cap (**Figure 6b**). This is normal.
434
435
436



437
438 **Figure 6a**

Figure 6b

439
440 c. Once the injection site is dry, remove the **yellow** inner needle cap (**Figure 6c**). You are
441 now ready to inject.
442



443

444 **Figure 6c**

445

446 **d. Hold the PegIntron REDIPEN with your fingers wrapped around the pen body**
 447 **barrel and your thumb on the dosing button (Figure 7).**

448

- With your other hand, pinch the skin in the area you have cleaned for injection.

449

- Insert the needle into the pinched skin at an angle of 45° to 90°.

450

- Press the dosing button down slowly and firmly until you can't push it any further.

451

- Keep your thumb pressed down on the dosing button for an additional 5 seconds to ensure that you get the complete dose.

452

453

- Remove the needle from your skin.



454

455 **Figure 7**

456 **e. Gently press the injection site with a small bandage or sterile gauze if necessary for a**
 457 **few seconds but do not massage the injection site. If there is bleeding, cover with an**
 458 **adhesive bandage. DO NOT RECAP THE NEEDLE and DO NOT REUSE the**
 459 **REDIPEN.**

460

461 **How do I Dispose of the REDIPEN?**

462 Discard the REDIPEN and needle and any solution remaining in the REDIPEN in a sharps
 463 container or other puncture-resistant container like a metal coffee can. DO NOT use glass or
 464 clear plastic containers. Ask your health care provider how to dispose of a full container.
 465 Always keep the container out of reach of children.

466

467 **After 2 hours, check the injection site for redness, swelling, or tenderness.**

468 **If you have a skin reaction and it doesn't clear up in a few days, contact**

469 **your health care provider.**

470

471 Manufactured by: Schering Corporation, Kenilworth, NJ 07033 USA

472 **DATE 12/08**

473 **27853560T**